

Maternal Depressive Symptoms Not Associated with Reduced Height in Young Children in a US Prospective Cohort Study

Karen A. Ertel^{1*}, Karestan C. Koenen^{1,2,3}, Janet W. Rich-Edwards^{2,4}, Matthew W. Gillman^{5,6}

1 Department of Society, Human Development, and Health & Kellogg Health Scholars Program, Harvard School of Public Health, Boston, Massachusetts, United States of America, **2** Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts, United States of America, **3** Center on the Developing Child, Harvard University, Cambridge, Massachusetts, United States of America, **4** Center for Women's Health and Gender Biology, Brigham and Women's Hospital, Boston, Massachusetts, United States of America, **5** Obesity Prevention Program, Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, Massachusetts, United States of America, **6** Department of Nutrition, Harvard School of Public Health, Boston, Massachusetts, United States of America

Abstract

Background: Shorter stature is associated with greater all cause and heart disease mortality, but taller stature with increased risk of cancer mortality. Though childhood environment is important in determining height, limited data address how maternal depression affects linear growth in children. We examined the relationships between antenatal and postpartum depressive symptoms and child height and linear growth from birth to age 3 years in a U.S. sample.

Methods: Subjects were 872 mother-child pairs in Project Viva, a prospective pre-birth cohort study. The study population is relatively advantaged with high levels of income and education and low risk of food insecurity. We assessed maternal depression at mid-pregnancy (mean 28 weeks' gestation) and 6 months postpartum with the Edinburgh Postnatal Depression Scale (score ≥ 13 on 0–30 scale indicating probable depression). Child outcomes at age 3 were height-for-age z-score (HAZ) and leg length. HAZ was also available at birth and ages 6 months, 1, 2, and 3 years.

Findings: Seventy (8.0%) women experienced antenatal depression and 64 (7.3%) experienced postpartum depression. The mean (SD) height for children age 3 was 97.2 cm (4.2), with leg length of 41.6 cm (2.6). In multivariable linear regression models, exposure to postpartum depression was associated with greater HAZ (0.37 [95% confidence interval: 0.16, 0.58]) and longer leg length (0.88 cm [0.35, 1.41]). The relationship between postpartum depression and greater HAZ was evident starting at 6 months and continued to age 3. We found minimal relationships between antenatal depression and child height outcomes.

Conclusion: Our findings do not support the hypothesis that maternal depression is associated with reduced height in children in this relatively advantaged sample in a high-income country.

Citation: Ertel KA, Koenen KC, Rich-Edwards JW, Gillman MW (2010) Maternal Depressive Symptoms Not Associated with Reduced Height in Young Children in a US Prospective Cohort Study. PLoS ONE 5(10): e13656. doi:10.1371/journal.pone.0013656

Editor: Landon Myer, University of Cape Town, South Africa

Received: May 31, 2010; **Accepted:** September 18, 2010; **Published:** October 27, 2010

Copyright: © 2010 Ertel et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Project Viva is supported by grants from the National Institutes of Health (HD 34568, HL 64925, HL 68041). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: kertel@hsph.harvard.edu

Introduction

Extant literature largely from the United States and Western European nations reveals that body height is strongly associated with health outcomes. Shorter stature is associated with increased risk of mortality,[1–5] as well as mortality due to cardiovascular[1,2] and respiratory disease.[1,4] At the other end of the spectrum, tall individuals have greater risk for cancer and mortality due to cancer.[4,6–9] The underlying reasons for these associations are unknown, but two possible mechanisms are that adult height reflects growth, nutrition, and the social environment in early life and that height in adulthood is positively associated with social class and its attendant exposures.[1] Leg length, a component of height, may be a more sensitive marker of prepubertal environmental influences than total height because

height increases in childhood are due more to leg growth than trunk growth.[10] Similar to total height, leg length is inversely associated with cardiovascular risk factors and coronary heart disease,[6,7,11–13] but positively associated with cancer risk.[6]

Height in childhood is strongly correlated with adult height: at age 2 years, length or height has a correlation of 0.8 with adult height.[14] Similar to attained height in adulthood, accelerated linear growth in childhood has been positively associated with breast cancer risk.[15,16] Additionally, a recent analysis of over 3,000 young girls in the United States revealed that greater height in childhood (ages 4–5 years) predicted early menarche,[17] which is linked to adverse mental and physical health outcomes.[15,18,19]

The current literature regarding determinants of height focuses on developing countries. This is critically important, as nutritional

status, disease, and infrastructure have significant influences on attained height in this context. However, the evidence reviewed above for the associations between height and adult health outcomes comes largely from the United States or Western European nations, thus we believe there is value in continued investigation of determinants of linear growth in the context of developed countries, such as the United States. Though many factors contribute to final height, including genetics, nutrition, and disease,[20] less attention has been paid to potential social and psychosocial influences on linear growth. Maternal mental health is a primary determinant of the child's psychosocial environment. Maternal depression is common in the postpartum period, with an estimated 14% of new mothers experiencing depression in the first 6 months after delivery.[21] Given its prevalence and its documented effect on child environment,[22,23] mother-child interaction,[24–26] and child physiology,[27–29] maternal depression may be an important determinant of linear growth in young children.

Previous studies using underweight or non-organic failure to thrive as indicators of poor physical growth have suggested an association between maternal depression and inadequate growth, but results have not consistently supported this association [30–37] (for a review of recent findings, see Stewart et al, 2007[38]). The literature specific to height is smaller and largely in developing countries; these studies provide mixed support for an association between postpartum depression and reduced child linear growth. Studies in India,[39] Bangladesh,[40] Brazil,[41] and Nigeria[42] have shown an association between maternal depressive symptoms and child stunting, while studies in South Africa[43] and Brazil[44] have shown no significant difference in child height according to maternal depressive symptoms. Similar methodology was used in four developing countries to examine maternal mental health (not depression per se) and child stunting; they found an independent, cross-sectional association between maternal 'common mental disorder' and stunting in India, but not Ethiopia, Peru, or Vietnam.[45] It is difficult to compare across studies due to different measures of maternal depression and ages of assessment of children. Further complicating interpretation is the possibility of reverse causation, as most of these studies were cross-sectional.

Antenatal depression, through its effects on fetal growth, may also influence child stature. Antenatal depression, with a period prevalence of 18% during pregnancy,[21] is associated with low birth weight, preterm birth, and small for gestational age at birth.[46] Infants born small tend to remain small in height as well as weight through childhood,[47] suggesting that an adverse intrauterine environment may contribute to short stature. A prospective cohort study in India reported 3.2 and 2.8 increased odds of stunting at 6 months and 12 months, respectively, for children whose mothers were depressed in the 3rd trimester of pregnancy compared to children whose mothers were not depressed at that time.[48]

Based on this prior literature, we hypothesized that children whose mothers experienced antenatal or postpartum depression would be shorter than children of mothers without depression. This study is among the first to examine these associations in a high-income country. Additionally, the sample population for this study differs from those in which an association has been found between maternal depression and stunting in several important ways, including higher levels of family income, access to sophisticated health services, low prevalence of food insecurity and hunger, low rates of infection, high levels of education, and low rates of both depression and stunting. Thus, this analysis may represent a rather stringent test of the association between maternal depression and child height.

Methods

Sample and data collection

Subjects were from Project Viva, a prospective cohort study of pregnant women and their singleton children. Participants were recruited at their first prenatal visit from eight obstetric practices of Harvard Vanguard Medical Associates, a large group practice in the greater Boston area. Exclusion criteria included multiple gestation, inability to answer questions in English, plans to move out of the area before delivery, and gestational age more than 22 weeks at first prenatal visit. 2670 pregnant women (64% of those eligible) were enrolled between April 1999 and July 2002; 329 subsequently became ineligible (60% because they were no longer pregnant). Of the remaining 2341 enrolled women, 195 withdrew and 18 were lost to follow-up before delivery. 2128 participants delivered a live infant and were eligible for inclusion in the current analysis. 1635 (70%) completed a mid-pregnancy questionnaire that assessed depressive symptoms and 1422 (60% of those enrolled) completed a second questionnaire that assessed depressive symptoms at 6 months postpartum. 1249 women (53% of those enrolled) had data on both antenatal and postpartum depression. Data were also collected at birth, including enrollment of the child, and at 1, 2, and 3 years after delivery. Among participants with depressive symptom data at mid-pregnancy and 6 months postpartum, 928 had height data for children at age 3 years. Of these, we excluded 56 (6.0%) due to missing covariate information for a final sample size of 872. Comparison of the 872 subjects included in this analysis to those excluded revealed some differences: included mothers were more likely to be white (79.6% vs. 74.3%), more highly educated (38.4% vs. 22.1% with a graduate degree), slightly older (mean age 33 years vs. 31 years), and slightly taller (mean height 65.1 vs. 64.7 inches). The Project Viva study and consent procedures were approved by human subjects committees of Harvard Pilgrim Health Care, Brigham and Women's Hospital, and Beth Israel Deaconess Medical Center. Written informed consent was obtained from all mothers in the study.

Measures

Depressive symptoms. We assessed depressive symptoms with the 10-item Edinburgh Postnatal Depression Scale (EPDS) in mid-pregnancy (mean of 28 weeks' gestation) and at approximately 6 months after birth. We chose the EPDS because it has been validated for antenatal and postpartum use.[49] We used the cutpoint of 13 or more (on the 0–30 point scale) to indicate probable depression, consistent with previous work in our cohort [50,51] and in other large cohorts that collected EPDS data antenatally and postnatally.[52] The cut-point at 13 or more indicates probable depression with a sensitivity of 86% and specificity of 78% in the postnatal period.[49] The optimal cut-off for probable major depression in the antenatal period may be higher (15 or more),[53,54] thus we report multivariable results using this alternate cut-off. It should be noted that the EPDS is a screening tool that measures probable depression, and is not a clinical diagnosis of depression; however, to be succinct, we refer to an EPDS score ≥ 13 as antenatal or postpartum depression.

Child anthropometric measurements. Child length or height was assessed at birth, 6 months, 1 year, 2 years, and 3 years of age. Project Viva trained research staff completed measurements at birth, 6 months, and 3 years. We obtained measurements for one and two years of age from clinical records. In a previous study of patients in this group practice, Rifas-Shiman and colleagues found that clinical assessment of length or height in children under age 2 years systematically underestimated height

compared to research standard methods, and they developed a regression equation to correct for this bias.[55] In the analyses presented here, we employed the corrected lengths for clinical measurements of children at 1 and 2 years. At age 3, staff also measured sitting height. We calculated leg length at age 3 as the difference between standing height and sitting height. Length/height was measured to the nearest 0.1 centimeter. Details regarding the age-three assessment have been published.[56] We calculated age- and sex-specific height-for-age z-score (HAZ) according to US national reference data for birth and ages six months, 1, 2, and 3 years.[57] We also calculated age- and sex-specific height-for-age z-score for age 3 years according to World Health Organization (WHO) Child Growth Standards[58], using the SAS program provided by WHO (available at <http://www.who.int/childgrowth/software/en/>, accessed 9/2/2010).

Sociodemographic factors and potential confounders. We chose variables that have been previously linked with perinatal depression or child height as covariates, see Table 1. Maternal age, race/ethnicity, household income, education, partnership status, height, pre-pregnancy BMI, and maternal report of the father's height were recorded at enrollment. We assessed maternal smoking during pregnancy by questionnaire and categorized responses into former smoker, never smoker, and smoked during pregnancy. We calculated pregnancy weight gain as pre-pregnancy weight subtracted from the last clinically recorded weight before delivery and categorized it according to the Institute of Medicine's gestational weight gain guidelines.[59] Presence of gestational diabetes or impaired glucose tolerance during pregnancy was collected from medical records. We identified women who used medication for depression during pregnancy by searching prescription data for the 90 days before the last menstrual period through delivery for 29 commonly used antidepressants. We calculated gestational age at delivery from the last menstrual period; if estimated gestational age based on last menstrual period differed by more than 10 days from the 2nd trimester ultrasound estimate, we used the estimate from the ultrasound. We calculated birthweight for gestational age z-scores with use of US national reference data.[60]

Potential mediators. We considered the following to be potential mediators of antenatal and/or postpartum depression's impact on child height: subsequent depression (at 6 months and 1 year for antenatal depression and at 1 year for postpartum depression, each assessed by the EPDS), duration of breastfeeding, and age at which solid foods were introduced to the child, all of which were assessed by questionnaire. There was little overlap between depressive symptoms occurring in the perinatal period with depressive symptoms at 1 year postpartum: 18 subjects had elevated depressive symptoms at both 6 months and 1 year postpartum and 11 at both antenatal and 1 year postpartum. Additionally, we considered gestational age at birth and birthweight for gestational age potential mediators of the influence of antenatal depression on child stature.

Statistical analysis

We examined two sets of outcomes: HAZ and leg length at age 3 years, and change in HAZ from birth through age 3. In both cases, we separately modeled the impact of antenatal (EPDS ≥ 13 and EPDS ≥ 15) and postpartum depression. For HAZ and leg length at age 3, we used multivariable linear regression models. In the base model (Model 1) for each exposure, we adjusted for child sex and age in months at assessment. In Model 2, we adjusted for sociodemographic and physiological factors related to maternal depression and/or child height. In Model 3, we additionally adjusted for gestational age at birth and birthweight for gestational

age z-score; these variables may be mediators of the association of antenatal depression with height as well as confounders of the association of postpartum depression with height. To increase precision of the estimate of interest, if removal of a covariate did not substantially alter the effect estimate of the primary exposure, that covariate was not included in the models presented here. Covariates tested and not included were maternal marital status, education, smoking during pregnancy, gestational diabetes or impaired glucose tolerance during pregnancy, paternal height, and anti-depressant use during pregnancy. Our primary outcomes were HAZ according to CDC reference standards and leg length; we also ran our models using HAZ according to WHO reference standards because the CDC and WHO standards are based on different reference populations. If results are consistent using both standards, we have some assurance that estimated associations are not due to the choice of reference population.

Seventy mothers experienced depression during pregnancy and 64 at 6 months postpartum. Only 19 mothers experienced depression at both time points. Due to the relative lack of overlap in these exposures, we examined antenatal and postpartum depression separately. Estimated effects for antenatal and postpartum depression were similar whether we included these 19 subjects or not; thus they are included in both exposure categories.

Research-quality measurements of birth length were available in a subset of the study population ($n = 516$). In this subset, we found that controlling for birth length z-score in place of birthweight for gestational age z-score (available for the whole cohort) did not affect results, so we present results for the larger cohort only, and use birthweight for gestational age z-score as a control variable. Also in this subset of the study population, we conducted a direct test of reverse causation for the association between postpartum depression and height outcomes at age 3: we examined whether length at birth predicted postpartum depression, controlling for maternal age, race/ethnicity, height, pregnancy weight gain, household income, and child gender.

Potential mediators of the effect of antenatal depression were gestational age at birth, birthweight for gestational age,[46] breastfeeding duration, age of introduction of solid foods, and maternal depression at 6 months and 1 year postpartum. Potential mediators of the association of postpartum depression with height were breastfeeding duration, age of introduction of solid foods, and depression at 1 year postpartum. To test for mediation, we added potential mediators to the fully-adjusted models and assessed the magnitude of attenuation of the estimated effect of exposure to maternal depression.

For the outcome change in HAZ from birth to 3 years, we used longitudinal models allowing a random intercept, random slope, and an unstructured covariance matrix. Covariates were based on the fully-adjusted models presented in Tables 2 and 3. In models with antenatal depression as primary predictor, we modeled the change in HAZ from birth through age 3. In models with postpartum depression as primary predictor, we modeled the change in HAZ from age 6 months to 3 years. We entered time as a categorical variable, with each outcome assessment (birth, 6 months, 1, 2, and 3 years) as a category and tested for an interaction between maternal depression and time. All analyses employed SAS version 9.1 (SAS Institute, Cary, NC), `proc glm` for age 3 outcomes and `proc mixed` for longitudinal analyses.

Results

Mothers in this sample were predominantly white, married, and had an annual household income over \$70,000 (Table 1). Seventy (8.0%) participants experienced antenatal depression and 64

Table 1. Characteristics of Project Viva Participants According to Maternal Antenatal and Postpartum Depression. Data from 872 mother-child pairs participating in Project Viva.

	Overall (n = 872)		Antenatal Depression				Postpartum Depression			
	mean/n	SD/%	No (n = 802)		Yes (n = 70)		No (n = 808)		Yes (n = 64)	
			mean/n	SD/%	mean/n	SD/%	mean/n	SD/%	mean/n	SD/%
Maternal characteristics										
Age	33.0	4.5	33.0	4.4	32.1	5.0	33.1	4.4	31.6	5.4
Race/ethnicity										
Black	73	8.4	64	8.0	9	12.9	65	8.0	8	12.5
Other	105	12.0	92	11.5	13	18.6	94	11.6	11	17.9
White	694	79.6	646	80.6	48	68.6	649	80.3	45	70.3
Education										
Less than BA/BS	189	21.7	169	21.1	20	28.6	169	20.9	20	31.3
BA/BS	348	39.9	322	40.2	26	37.1	327	40.5	21	32.8
Graduate degree	335	38.4	311	38.8	24	34.3	312	38.6	23	35.9
Married *	784	90.0	728	90.9	56	80.0	733	90.8	51	79.7
Household income										
< = \$40,000	77	8.8	61	7.6	16	22.9	63	7.8	14	21.9
40,001 to \$70,000	195	22.4	176	22.0	19	27.1	176	21.8	19	26.7
> = \$70,001	600	68.8	565	70.5	35	50.0	569	70.4	31	48.4
Pre-pregnancy BMI > = 25	288	33.0	259	32.3	29	41.4	260	32.2	28	43.8
Height (inches)	65.1	2.7	65.1	2.7	64.6	2.4	65.1	2.7	64.8	2.9
Paternal height (inches)	70.7	3.0	70.7	3.0	70.6	3.0	70.7	2.9	70.5	3.3
BF duration (months) *	6.55	4.5	6.50	4.5	7.11	4.7	6.57	4.5	6.31	4.3
Introduction of solid foods *										
before 4 months	130	14.9	125	15.6	5	7.1	119	14.8	11	17.2
4–5 months	623	71.5	566	70.7	57	81.4	578	71.6	45	70.3
6 months or after	118	13.6	110	13.7	8	11.4	110	13.6	8	12.5
Infant characteristics										
Girl	456	52.3	417	52.0	39	55.7	427	52.9	29	45.3
GA at delivery (weeks)	39.6	1.7	39.6	1.7	39.5	1.5	39.6	1.7	39.5	1.6
BW for GA (z value)	0.23	0.93	0.25	0.93	0.04	0.95	0.24	0.92	0.13	1.07
Child characteristics at 3 yrs										
Age (months)	39.0	3.2	39.0	3.2	38.8	2.8	39.0	3.2	38.6	2.6
Height (cm)	97.2	4.2	97.2	4.2	97.1	4.6	97.1	4.2	98.6	4.0
HAZ	0.2	0.9	0.2	0.9	0.2	1.0	0.2	0.9	0.6	0.9
Leg Length (cm)	41.6	2.6	41.6	2.5	42.0	2.8	41.6	2.6	42.4	2.3

BA/BS, Bachelor of Arts or Science; BMI, body mass index (kg/m²); BF, breastfeeding; GA, gestational age; BW, birth weight; HAZ, height-for-age z-score.

*Sample sizes are slightly different due to missing data: breastfeeding duration n = 869; marital status n = 871; introduction of solid foods n = 871.

doi:10.1371/journal.pone.0013656.t001

(7.3%) experienced postpartum depression. Women with antenatal depression were less likely to be married and more likely to have lower income, inadequate weight gain during pregnancy, and gestational diabetes or impaired glucose tolerance. Women with postpartum depression were younger, less likely to be married, had lower income, and their children were taller and had longer leg length at age 3.

In multivariable models predicting age 3 outcomes, there was some indication that exposure to antenatal depression was associated with greater leg length but not HAZ (Table 2). In a model controlling for potential confounders (Model 2), antenatal depression (EPDS > = 13) was associated with (0.45 (−0.07, 0.97)) greater leg length. Use of the more stringent definition of antenatal

depression (EPDS > = 15) revealed a stronger association between antenatal depression and leg length in children (0.74 (0.07, 1.40)). Further control for depression at 6 months and 1 year postpartum did not materially alter results.

Postpartum depression was associated with greater HAZ and leg length. As Table 3 displays, after controlling for potential confounders (Model 3), compared to children not exposed to postpartum depression, exposed children had HAZ 0.37 (0.16, 0.58) and leg length 0.88 cm (0.35, 1.41) larger. Maternal height, birthweight for gestational age, non-white race/ethnicity, and lower household income were other factors associated with taller height in children (Table 4). The finding that black children of this age are taller is consistent with findings from a nationally-

Table 2. Difference in 3-year height outcomes among children exposed versus not exposed to antenatal depression, using 2 alternate cut-off values for the Edinburgh Postnatal Depression Scale (EPDS).

	EPDS >= 13		EPDS >= 15	
	Estimate (95% confidence interval)		Estimate (95% confidence interval)	
	Height-for-age z-score	Leg Length (cm)	Height-for-age z-score	Leg Length (cm)
Model 1	0.002 (-0.22, 0.23)	0.48 (-0.08, 1.04)	0.07 (-0.21, 0.36)	0.63 (-0.08, 1.34)
Model 2	0.01 (-0.20, 0.22)	0.45 (-0.07, 0.97)	0.13 (-0.14, 0.39)	0.74 (0.07, 1.40)
Model 3	0.03 (-0.17, 0.24)	0.48 (-0.03, 1.00)	0.16 (-0.10, 0.42)	0.79 (0.14, 1.44)
Model 4	-0.04 (-0.25, 0.17)	0.32 (-0.20, 0.84)	0.06 (-0.21, 0.32)	0.58 (-0.09, 1.24)
Model 5	-0.02 (-0.23, 0.19)	0.36 (-0.16, 0.89)	0.10 (-0.17, 0.37)	0.68 (0.01, 1.35)

Model 1 covariates: child sex and age at 3-year assessment.

Model 2: Model 1+ maternal: age, race/ethnicity, household income, height, pregnancy weight gain.

Model 3: Model 2+ gestational age at birth and birthweight for gestational age z-value.

Model 4: Model 3+ postpartum depression.

Model 5: Model 4+ breastfeeding duration and age of introduction of solid foods (n=869).

Data from 872 mother-child pairs participating in Project Viva.

doi:10.1371/journal.pone.0013656.t002

representative sample of young children.[61] Restriction to individuals without antenatal depression did not materially alter results, nor did further control for maternal depression at 1 year postpartum. There was no evidence for mediation by breastfeeding duration or age of introduction of solid foods, though longer duration of breastfeeding was an independent predictor of reduced HAZ and leg length. Each month of longer duration of breastfeeding was associated with a -0.02 (-0.03 , -0.01) difference in HAZ and a -0.05 cm (-0.08 , -0.01) difference in leg length. Additionally, there was no evidence of reverse causation: birth length was not a significant predictor of postpartum depression (OR = 1.39, 95% CI: 0.85, 2.27) in a logistic regression model controlling for maternal age, race, height, pregnancy weight gain, household income, and child gender.

Use of WHO reference standards revealed similar results as those displayed in Table 2 and Table 3. The estimated association between antenatal depression and WHO height-for-age z-score was -0.01 (-0.27 , 0.24) in a model controlling for maternal age, race/ethnicity, household income, height, and pre-pregnancy weight gain (the same covariates as in Model 2 of Table 2). The

estimated association between postpartum depression and WHO HAZ was 0.28 (0.02 , 0.54) controlling for the same covariates as in Model 3 of Table 3.

In longitudinal models, the association between postpartum depression and child HAZ from 6 months through age 3 was similar for each age (Figure 1), thus we did not include a postpartum depression by age interaction term in the final model. The association between postpartum depression and child HAZ was 0.29 (0.11 , 0.47) at each time point. Similar to age 3 findings, antenatal depression was not associated with HAZ from birth through age 3: the estimated association over all time points was -0.01 (-0.19 , 0.17).

Discussion

This study did not find evidence for the hypothesis that maternal depression is associated with reduced height in children. Contrary to expectations, maternal postpartum depression was associated with greater child total height and leg length in this relatively privileged sample in a high income country. Postpartum depression was associated with 0.37 higher HAZ, which translates into an approximately 1.5 centimeter difference in 3-year-olds of average height (94.5 cm) in the US.[57] This finding was independent of potential confounders, including maternal socio-demographic factors and height as well as child size at birth. We found no evidence for mediation by duration of breastfeeding or age of introduction of solid foods, nor any evidence for reverse causation. Additional control for depression at 1 year postpartum did not diminish the estimated effect for postpartum depression at 6 months, nor was there a main effect of depressive symptoms at 1 year postpartum on height outcomes, suggesting that exposure to maternal depression earlier in life is more important.

Antenatal depression when defined by the more stringent criteria of EPDS >= 15 was associated with greater leg length in children. Comparing results using alternate cut-off values for the EPDS suggests a dose-response relation between antenatal depressive symptoms and linear growth outcomes in children.

Our results contrast with several studies in developing countries that report either a cross-sectional or prospective association between maternal depressive symptoms or clinical depression and reduced child height or increased risk of stunting.[39–42] However, not all studies in developing countries support an association

Table 3. Difference in 3-year height outcomes among children exposed versus not exposed to postpartum depression.

	Estimate (95% confidence interval)	
	Height-for-age z-score	Leg Length (cm)
Model 1	0.39 (0.16, 0.62)	0.97 (0.39, 1.56)
Model 2	0.35 (0.13, 0.57)	0.85 (0.31, 1.39)
Model 3	0.37 (0.16, 0.58)	0.88 (0.35, 1.41)
Model 4	0.37 (0.16, 0.58)	0.89 (0.36, 1.41)

Model 1 covariates: child sex and age at 3-year assessment.

Model 2: Model 1+ maternal: age, race/ethnicity, household income, height, pregnancy weight gain.

Model 3: Model 2+ gestational age at birth and birthweight for gestational age z-value.

Model 4: Model 3+ breastfeeding duration and age of introduction of solid foods (n=869).

Data from 872 mother-child pairs participating in Project Viva.

doi:10.1371/journal.pone.0013656.t003

Table 4. Factors associated with height outcomes at age 3 years.

Variable	Estimate (95% confidence interval)	
	Height-for-age z-score	Leg Length (cm)
Postpartum depression	0.37 (0.16, 0.58)	0.88 (0.35, 1.41)
Maternal race/ethnicity		
Black	0.39 (0.19, 0.60)	2.04 (1.53, 2.56)
Other	0.18 (0.00, 0.35)	0.57 (0.14, 1.01)
White	0 (ref)	0 (ref)
Household income		
<= \$40,000	0.20 (0.00, 0.41)	0.33 (-0.19, 0.85)
40,001 to \$70,000	0.12 (-0.02, 0.25)	0.16 (-0.17, 0.50)
>= \$70,001	0 (ref)	0 (ref)
Maternal height (inches)	0.11 (0.09, 0.13)	0.24 (0.19, 0.29)
Birthweight for gestational age z-score	0.23 (0.17, 0.29)	0.35 (0.20, 0.51)

Multivariable linear regression model adjusted for maternal age and pregnancy weight gain and child sex, gestational age at birth, and age at outcome assessment as well as all variable in the Table.
doi:10.1371/journal.pone.0013656.t004

between maternal mental health and reduced height in children.[43–45] It should also be noted that maternal depression has been found to be associated with both reduced weight[32,34,39,62] (for a review, see Stewart, 2007 [38]) and *increased* weight[63] in different settings; thus, there is prior evidence that maternal depression may have different effects in different contexts.

It is unclear what mechanisms may explain divergent findings between developing countries and the US. One possible source of difference is risk of nutritional deficiency and serious childhood illness across these settings. In resource-poor settings, maternal depression may be associated with or lead to lack of sufficient food for the household or inadequate care of childhood illness, both of which may contribute to shorter stature in children. In Project Viva, a middle and upper class sample in the US, there are likely very few families who cannot obtain adequate food, though we do not have data to examine this issue directly. Similarly, severe childhood illness that may affect height is limited in the Project Viva sample. Other notable differences between the study population examined here and studies in low and middle income countries include high levels of maternal education, relatively low levels of maternal depression, and low rates of child stunting. We may speculate that in the rather privileged families enrolled in Project Viva, one threat (maternal depressive symptoms) is not enough to interrupt children’s growth, while in developing countries in the context of several threats (including food insecurity, low education, and high rates of stunting) and fewer resources, maternal depression is an additional burden that influences children’s growth.

Another possible reason for the different findings is that many of the studies in developing countries used a dichotomous outcome of stunting, while we used height as a continuous outcome. We cannot examine whether this may help explain divergent results because only 7 subjects in our sample met criteria for stunting (HAZ<-2). Reports from Brazil, however, support the notion that maternal depression is associated with stunting,[41] but not height when examined as a linear outcome variable.[44] Future research may endeavor to examine these possible explanations further.




Figure 1. Estimated longitudinal association of postpartum depression with child height-for-age z-score (HAZ). Data from 872 mother-infant pairs participating in Project Viva. Multivariable regression estimates adjusted for maternal: age, race/ethnicity, household income, height, and weight gain during pregnancy and child: sex, gestational age, and birthweight for gestational age z-score. Bars indicate 95% confidence intervals. Exposed to postpartum depression (dotted line), not exposed to postpartum depression (solid line).
doi:10.1371/journal.pone.0013656.g001

Interpretation of our results should be in light of the study's limitations. Mother-child pairs included in our analysis represent 37% of enrolled women. It is possible that this loss to follow-up introduced some bias, though it is unlikely that it induced a positive association between postpartum depression and child height. This loss to follow-up may limit the generalizability of our findings. The mothers included are generally well-educated, of middle to upper income, and all have health insurance. It is possible that the association between maternal depression and child height may be different in more at-risk populations. Similarly, though we statistically control for important variables related to maternal depression and height, residual confounding is always a concern. However, while residual confounding could have resulted in over- or under-estimates in our study, it is unlikely that the direction of effect would have changed with better control of confounding. Regarding our exposures, we rely on self-report of depressive symptoms, not clinically-defined depression, and depressive symptoms were assessed at only two time points in the perinatal period, which likely resulted in misclassification of exposure. This may be particularly important for postpartum depression: we assessed postpartum depression at 6 months, but the prevalence of postpartum depression is highest in the 2nd and 3rd months postpartum.[21] Misclassification of our exposure is likely nondifferential with respect to the outcomes, which would attenuate results towards the null. Finally, due to the relative independence of antenatal and postpartum depression, our results generalize primarily to women who have either antenatal or postpartum depression, although inclusion of women with both did not materially alter the estimated effects in this cohort of women. Given these limitations as well as the unexpected findings, replication of these results in another US-based cohort is particularly important.

In considering the mechanisms underlying the current findings, although many factors influence height, very few are likely to mediate an effect of postpartum depression on linear growth. One possible pathway is suboptimal nutritional intake among children of depressed mothers. Depressed mothers may have difficulty recognizing signs of satiety in their children, use more mechanistic methods of feeding, or use restrictive feeding practices.[64,65] Secondly, maternal depression negatively affects mother-child interactions,[24,25] which may have implications for the child's nutritional intake. Depressed mothers may have more unhealthy behaviors, such as dysregulated eating behaviors (including over-eating and eating high-calorie foods) and limited physical activity, and their children may have similar behavior patterns. It is possible that these behaviors lead to increased energy intake in infants; in a related analysis of Project Viva participants, maternal postpartum depression was associated with greater overall adiposity in children at age 3 years.[66] We may speculate that increased caloric intake could have contributed to the greater linear growth observed here through actions of growth hormone or insulin-like growth factors; however, this study did not measure caloric intake or growth hormone, thus we were not able to test

these possible mechanisms. That children exposed to postpartum depression showed greater height as early as 6 months of age suggest that influences before this age are likely to be at play.

Another possible pathway is through cortisol. In two studies, postpartum depression was associated with increased cortisol and cortisol variability in newborns and adolescents.[27,67] Postpartum depression, through its adverse effects on parenting behavior and the social environment, can produce more stressful experiences for the child. Stressful experiences, in turn, lead to elevated cortisol levels and increased cortisol reactivity in children.[68,69] Cortisol has a complex relationship with growth hormone: while chronic hypercortisolism leads to reduced growth hormone and reduced growth, small increases in cortisol stimulate growth hormone secretion and production.[70] It is possible that increased cortisol in the children exposed to postpartum depression had a stimulating effect on growth hormone and thereby increased height. Unfortunately, we did not measure these hormones in the current study.

The current study has several strengths. Project Viva is a prospective study with repeated outcome measurements, which allowed us to establish temporal order between exposure and outcomes as well as examine change over time in outcomes. This study assessed depressive symptoms both during pregnancy and during the postpartum period, allowing us to assess the contribution of each exposure individually. In addition, the primary outcomes of this study were collected using research-standard protocols, improving the precision with which we estimate relations.

Our study did not support the hypothesis that maternal depression is associated with shorter stature in children. To our knowledge, this study is the first to report an association between maternal depression and increased height in children. As one of the first studies to examine these relationships in a US sample, these findings should be replicated in other, more diverse samples in the US and other developed nations. Our findings suggest that postpartum depression has important influences on behavior or physiology early in life that translate into different growth patterns in children. Public health implications of our findings await elucidation of the contexts in which and mechanisms through which maternal depression may affect growth in children. Future research should study possible mechanisms that may account for the different linear growth seen in children of depressed mothers, including mother-infant interactions, dysregulated feeding and/or eating behavior, as well as underlying hormonal differences.

Author Contributions

Analyzed the data: KAE. Wrote the paper: KAE. Conceived the research question: KAE. Interpreted data results: KAE KK JR-E MWG. Contributed to study concept: KK MWG. Revised article for intellectual content: KK JR-E MWG. Approved final version of article: KK JR-E MWG. Contributed to data acquisition: JR-E. Primary acquisition of data: MWG.

References

- Davey Smith G, Hart C, Upton M, Hole D, Gillis C, et al. (2000) Height and risk of death among men and women: aetiological implications of associations with cardiorespiratory disease and cancer mortality. *Journal of Epidemiology and Community Health* 54: 97–103.
- Jousilahti P, Tuomilehto J, Vartiainen E, Eriksson J, Puska P (2000) Relation of adult height to cause-specific and total mortality: a prospective follow-up study of 31,199 middle-aged men and women in Finland. *American Journal of Epidemiology* 151: 1112–1120.
- Peck A, Vagero DH (1989) Adult body height, self perceived health and mortality in the Swedish population. *Journal of Epidemiology and Community Health* 43: 380–384.
- Song Y, Sung J (2008) Adult height and the risk of mortality in South Korean women. *American Journal of Epidemiology* 168: 497–505.
- Waaler H (1984) Height, weight and mortality: the Norwegian experience. *Acta Medica Scandinavica* 679: 1–56.
- Gunnell D, Davey Smith G, Frankel S, Nanchahal K, Braddon FEM, et al. (1998) Childhood leg length and adult mortality: follow up of the Carnegie (Boyd Orr) Survey of diet and health in pre-war Britain. *Journal of Epidemiology and Community Health* 52: 142–152.
- Gunnell D, Okasha M, Davey Smith G, Oliver SE, Sandhu J, et al. (2001) Height, leg length, and cancer risk: A systematic review. *Epidemiologic Reviews* 23: 313–342.

8. Hebert P, Ajani U, Cook NR, Lee I, Chan KS, et al. (1997) Adult height and incidence of cancer in male physicians (United States). *Cancer Causes and Control* 8: 591–597.
9. Albanes D, Jones DY, Schatzkin A, Micozzi MS, Taylor PR (1988) Adult stature and risk of cancer. *Cancer Research* 48: 1658–1662.
10. Hall J, Allanson JE, Gripp KW, Slavotinek AM (2007) *Handbook of Physical Measurements*. Oxford UK: Oxford University Press.
11. Gunnell D, Whitley E, Upton MN, McConnachie A, Davey Smith G, et al. (2003) Associations of height, leg length, and lung function with cardiovascular risk factors in the Midspan Family Study. *Journal of Epidemiology and Community Health* 57: 141–146.
12. Lawlor D, Ebrahim S, Davey Smith G (2002) The association between components of adult height and Type II diabetes and insulin resistance: British Women's Heart and Health Study. *Diabetologia* 45: 1097–1106.
13. Lawlor D, Taylor M, Davey Smith G, Gunnell D, Ebrahim S (2004) Associations of components of adult height with coronary heart disease in postmenopausal women: the British women's heart and health study. *Heart* 90: 745–749.
14. Tanner J (1990) *Fetus into man: Physical growth from conception to maturity*. Cambridge MA: Harvard University Press.
15. De Stavola B, dos Santos Silva L, McCormack V, Hardy RJ, Kuh DJ, et al. (2004) Childhood growth and breast cancer. *American Journal of Epidemiology* 159: 671–682.
16. Hilakivi-Clarke L, Forsen T, Eriksson JG, Luoto R, Tuomilehto J, et al. (2001) Tallness and overweight during childhood have opposing effects on breast cancer risk. *British Journal of Cancer* 85: 1680–1684.
17. Salsberry P, Reagan PB, Pajer K (2009) Growth differences by age of menarche in African American and white girls. *Nursing Research* 58: 382–390.
18. Kautiiala-Heino R, Kosunen E, Rimela M (2003) Pubertal timing, sexual behavior and self-reported depression in middle adolescence. *Journal of Adolescence* 26: 531–545.
19. Remsburg K, Demerath EW, Schubert CM, Chumlea WC, Sun SS, et al. (2005) Early menarche and the development of cardiovascular disease risk factors in adolescent girls: The Fels Longitudinal Study. *Journal of Clinical Endocrinology and Metabolism* 90: 2718–2724.
20. Silventoinen K (2003) Determinants of variation in adult body height. *Journal of Biosocial Science* 35: 263–285.
21. Gavin N, Gaynes B, Lohr K, Meltzer-Brody S, Gartlehner G, et al. (2005) Perinatal depression: A systematic review of prevalence and incidence. *Obstetrics & Gynecology* 106: 1071–1083.
22. Burke L (2003) The impact of maternal depression on familial relationships. *International Review of Psychiatry* 15: 243–255.
23. McLearn K, Minkovitz CS, Strobino DM, Marks E, Hou W (2006) The timing of maternal depressive symptoms and mothers' parenting practices with young children: Implications for pediatric practice. *Pediatrics* 118: e174–e182.
24. Lovejoy M, Graczyk PA, O'Hare E, Neuman G (2000) Maternal depression and parenting behavior: a meta-analytic review. *Clinical Psychology Review* 20: 561–592.
25. Murray L, Cooper P, Hipwell A (2003) Mental health of parents caring for infants. *Archives of Women's Mental Health* 6: s71–s77.
26. Reck C, Hunt A, Fuchs T, Weiss R, Noon A, et al. (2003) Interactive regulation of affect in postpartum depressed mothers and their infants. *Psychopathology* 37: 272–280.
27. Bugental D, Martorell GA, Barraza V (2003) The hormonal costs of subtle forms of infant maltreatment. *Hormones and Behavior* 43: 237–244.
28. Dawson G, Frey K, Panagiotides H, Yamada E, Hessl D, et al. (1999) Infants of depressed mothers exhibit atypical frontal electrical brain activity during interactions with mother and with a familiar, non-depressed adult. *Child Development* 70: 1058–1066.
29. Martins C, Gaffan EA (2000) Effects of early maternal depression on patterns of infant-mother attachment: A meta-analytic investigation. *Journal of Child Psychology and Psychiatry* 41: 737–746.
30. Evans S, Reinhart VB, Succop RA (1972) Failure to thrive: A study of 45 children and their families. *Journal of the American Academy of Child Psychiatry* 11: 440.
31. Breunlin D, Desai VJ, Stone ME, Swilley JA (1983) Failure-to-thrive with no organic etiology. A critical review of the literature. *International Journal of Eating Disorders* 2: 25–49.
32. O'Brien L, Heycock EG, Hanna M, Jones PW, Cox JL (2004) Postnatal depression and faltering growth: A community study. *Pediatrics* 113: 1242–1247.
33. Drewett R, Blair P, Emmett P, Emond A, ALSPAC Study Team, (2004) Failure to thrive in the term and preterm infants of mothers depressed in the postnatal period: a population-based birth cohort study. *Journal of Child Psychology and Psychiatry* 45: 359–366.
34. Wright C, Parkinson KN, Drewett RF (2006) The influence of maternal socioeconomic and emotional factors on infant weight gain and weight faltering (failure to thrive): data from a prospective birth cohort. *Archives of Disease in Childhood* 91: 312–317.
35. O'Callaghan M, Hull D (1978) Failure to thrive or failure to rear? *Archives of Disease in Childhood* 53: 788–793.
36. Ramsay M, Gisel EG, McCusker J, Bellavance F, Platt R (2002) Infant sucking ability, non-organic failure to thrive, maternal characteristics, and feeding practices: a prospective cohort study. *Developmental Medicine & Child Neurology* 44: 405–414.
37. Singer L, Song L, Hill BP, Jaffe AC (1990) Stress and depression in mothers of failure-to-thrive children. *Journal of Pediatric Psychology* 15: 711–720.
38. Stewart R (2007) Maternal depression and infant growth - a review of recent evidence. *Maternal and Child Nutrition* 3: 94–107.
39. Patel V, DeSouza N, Rodrigues M (2003) Postnatal depression and infant growth and development in low income countries: a cohort study from Goa, India. *Archives of Disease in Childhood* 88: 34–37.
40. Black M, Baqui AH, Zaman K, Arifeen SE, Black RE (2009) Maternal depressive symptoms and infant growth in rural Bangladesh. *American Journal of Clinical Nutrition* 89(suppl): 951S–957S.
41. Surkan P, Kawachi I, Ryan LM, Berkman LF, Carvalho Vieira LM, et al. (2008) Maternal depressive symptoms, parenting self-efficacy, and child growth. *American Journal of Public Health* 98: 125–132.
42. Adewuya A, Ola BO, Aloba OO, Mapayi BM, Okeniyi JAO (2008) Impact of postnatal depression on infants' growth in Nigeria. *Journal of Affective Disorders* 108: 191–193.
43. Tomlinson M, Cooper PJ, Stein A, Swarz L, Moltano C (2006) Post-partum depression and infant growth in a South African peri-urban settlement. *Child Care Health and Development* 32: 81–86.
44. Surkan P, Ryan LM, Carvalho Vieira LM, Berkman LF, Peterson KE (2007) Maternal social and psychological conditions and physical growth in low-income children in Piaui, Northeast Brazil. *Social Science & Medicine* 64: 375–388.
45. Harpham T, Huttly T, De Silva MJ, Abramsky T (2005) Maternal mental health and child nutritional status in four developing countries. *Journal of Epidemiology and Community Health* 59: 1060–1064.
46. Field T, Diego M, Hernandez-Reif M (2006) Prenatal depression effects on the fetus and newborn: a review. *Infant Behavior & Development* 29: 445–455.
47. Binkin N, Yip R, Fleshood L, Trowbridge FL (1988) Birth weight and childhood growth. *Pediatrics* 82: 828–834.
48. Rahman A, Iqbal Z, Bunn J, Lovel H, Harrington R (2004) Impact of maternal depression on infant nutritional status and illness - A cohort study. *Archives of General Psychiatry* 61: 946–952.
49. Cox J, Holden HM, Sagovsky R (1987) Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry* 150: 782–786.
50. Rich-Edwards J, Mohlajee AP, Kleinman K, Hacker MR, Majzoub J, et al. (2008) Elevated mid-pregnancy corticotrophin-releasing hormone is associated with prenatal, but not postpartum, maternal depression. *Journal of Clinical Endocrinology and Metabolism* 93: 1946–1951.
51. Rich-Edwards J, Kleinman K, Abrams A, Harlow BL, McLaughlin TJ, et al. (2006) Sociodemographic predictors of antenatal and postpartum depressive symptoms among women in a medical group practice. *Journal of Epidemiology and Community Health* 60: 221–227.
52. Heron J, O'Connor TG, Evans J, Golding J, Glover V, the ALSPAC Study Team (2004) The course of anxiety and depression through pregnancy and the postpartum in a community sample. *Journal of Affective Disorders* 80: 65–73.
53. Murray D, Cox JL (1990) Screening for depression during pregnancy with the Edinburgh Depression Scale (EPDS). *Journal of Reproductive and Infant Psychology* 8: 99–107.
54. Matthey S, Henshaw C, Elliott, S, Barnett, B (2006) Variability in use of cut-off scores and formats on the Edinburgh Postnatal Depression Scale - implications for clinical and research practice. *Archives of Women's Mental Health* 9: 309–315.
55. Rifas-Shiman S, Rich-Edwards JW, Scanlon KS, Kleinman KP, Gillman MW (2005) Misdiagnosis of overweight and underweight in children younger than 2 years of age due to length measurement bias. *Medscape General Medicine* 7: 55.
56. Gillman M, Rich-Edwards JW, Huh S, Majzoub JA, Oken E, et al. (2006) Maternal corticotropin-releasing hormone levels during pregnancy and offspring adiposity. *Obesity* 14: 1647–1653.
57. National Center for Health Statistics (2000) *CDC Growth Charts*, United States.
58. WHO Multicentre Growth Reference Study Group (2006) *WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development*. Geneva: World Health Organization.
59. Institute of Medicine (1990) *Nutrition during pregnancy*. Washington DC: National Academy Press.
60. Oken E, Kleinman KP, Rich-Edwards JW, Gillman MW (2003) A nearly continuous measure of birth weight for gestational age using a United States national reference. *BMC Pediatrics* 3: 6.
61. Overpeck M, Hediger ML, Ruan WJ, Davis WW, Maurer KR, et al. (2000) Stature, weight, and body mass among young US children born at term with appropriate birth weights. *The Journal of Pediatrics* 137: 205–213.
62. Anoop S, Saravanan B, Joseph A, Cherian A, Jacob KS (2004) Maternal depression and low maternal intelligence as risk factors for malnutrition in children: a community based case-control study from South India. *Archives of Disease in Childhood* 89: 325–329.
63. Surkan P, Kawachi I, Peterson KE (2008) Childhood overweight and maternal depressive symptoms. *Journal of Epidemiology and Community Health* 62: e11.
64. Farrow C, Blissett JM (2005) Is maternal psychopathology related to obesogenic feeding practices at 1 year? *Obesity Research* 13: 1999–2005.

65. Hurley K, Black MM, Papas MA, Caulfield LE (2008) Maternal symptoms of stress, depression, and anxiety are related to nonresponsive feeding styles in a statewide sample of WIC participants. *The Journal of Nutrition* 138: 799–805.
66. Ertel K, Koenen KC, Rich-Edwards JW, Gillman MW (2010) Antenatal and postpartum depressive symptoms are differentially associated with early childhood weight and adiposity. *Paediatric and Perinatal Epidemiology* 24: 179–189.
67. Halligan S, Herbert J, Goodyer IM, Murray L (2004) Exposure to postnatal depression predicts elevated cortisol in adolescent offspring. *Biological Psychiatry* 55: 376–381.
68. Azar R, Paquette D, Zoccolillo M, Baltzer F, Tremblay RE (2007) The association of major depression, conduct disorder, and maternal overcontrol with a failure to show a cortisol buffered response in 4-month-old infants of teenaged mothers. *Biological Psychiatry* 62: 573–579.
69. Waynforth D (2007) The influence of parent-infant cosleeping, nursing, and childcare on cortisol and SIgA immunity in a sample of British children. *Developmental psychobiology* 46: 640–648.
70. Stratakis C (2006) Cortisol and growth hormone: clinical implications of a complex, dynamic relationship. *Pediatric Endocrinology Reviews* 3: 333–338.