

RESEARCH ARTICLE

Leukocyte Subtype Counts and Its Association with Vascular Structure and Function in Adults with Intermediate Cardiovascular Risk. MARK Study



click for updates

OPEN ACCESS

Citation: Gomez-Sanchez L, García-Ortiz L, Recio-Rodríguez JI, Patino-Alonso MC, Agudo-Conde C, Rigo F, et al. (2015) Leukocyte Subtype Counts and Its Association with Vascular Structure and Function in Adults with Intermediate Cardiovascular Risk. MARK Study. PLoS ONE 10(4): e0119963. doi:10.1371/journal.pone.0119963

Academic Editor: Ingo Ahrens, University Hospital Medical Centre, GERMANY

Received: August 3, 2014

Accepted: December 17, 2014

Published: April 17, 2015

Copyright: © 2015 Gomez-Sánchez et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by grants funded by the Spanish Ministry of Science and Innovation (MICINN) and Carlos III Health Institute/European Regional Development Fund (ERDF) (MICINN, ISCIII/FEDER) (Red RedIAPP RD06/0018, Research Groups: RD06/0018/2009, RD06/0018/0045/RD06/0018/0027) and by the Health Research Fund(FIS PI10/01088, PI10/02077, PI10/02043) and Regional Health Management of the Castilla and León(GRS

Leticia Gomez-Sánchez¹, Luis García-Ortiz^{1,2,3}, José I. Recio-Rodríguez^{1,2}, María C. Patino-Alonso^{1,2,4}, Cristina Agudo-Conde^{1,2}, Fernando Rigo⁵, Rafel Ramos^{6,7,8}, Ruth Martí^{6,8}, Manuel A. Gomez-Marcos^{1,2,3*}, for the MARK Group⁹

1 Primary Care Research Unit, The Alamedilla Health Center, Salamanca, Spain, **2** Castilla and León Health Service (SACyL), Biomedical Research Institute of Salamanca (IBSAL), Salamanca, Spain, **3** Medicine Department, University of Salamanca, Salamanca, Spain, **4** Statistics Department, University of Salamanca, Salamanca, Spain, **5** San Agustín Health Center, Isles Baleares Health Service (IBSALUT), Palma de Mallorca, Spain, **6** Research Unit Family Medicine, Jordi Gol Institute for Primary Care Research (IDIAP Jordi Gol), Girona, Spain, **7** Translab Research Group, Department of Medical Sciences, School of Medicine, University of Girona, Girona, Spain, **8** Girona Biomedical Research Institute (IDIBGI), Dr. Trueta University Hospital, Girona, Spain, **9** MARK Group, Research Network on Preventive Activities and Health Promotion (redIAPP), Girona, Spain

* magomez@usal.es

Abstract

Objectives

We investigated the relationship between leukocyte subtype counts and vascular structure and function based on carotid intima-media thickness, pulse wave velocity, central augmentation index and cardio-ankle vascular index by gender in intermediate cardiovascular risk patients.

Methods

This study analyzed 500 subjects who were included in the **MARK** study, aged 35 to 74 years (mean: 60.3 ± 8.4), 45.6% women. Measurement: Brachial ankle Pulse Wave Velocity (ba-PWV) estimate by equation, Cardio-AnkleVascular Index (CAVI) using the VaSera device and Carotid ultrasound was used to measure carotid Intima Media Thickness (IMT). The Mobil-O-Graph was used to measure the Central Augmentation Index (CAIx).

Results

Total leukocyte, neutrophil and monocyte counts were positively correlated with IMT ($p < 0.01$) in men. Monocyte count was positively correlated with CAIx in women ($p < 0.01$). In a multiple linear regression analysis, the IMT mean maintained a positive association with the neutrophil count ($\beta = 1.500$, $p = 0.007$) in men. CAIx maintained a positive association with the monocyte count ($\beta = 2.445$, $p = 0.022$) in women.

635/A/11). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Conclusion

The results of this study suggest that the relationship between subtype circulating leukocyte counts and vascular structure and function, although small, may be different by gender. In men, the neutrophil count was positively correlated with IMT and in women, the monocyte count with CAIx, in a large sample of intermediate-risk patients. These associations were maintained after adjusting for age and other confounders.

Trial Registration

ClinicalTrials.gov [NCT01428934](#)

Introduction

Atherosclerosis is, at least in part, an inflammatory process involving the infiltration of the circulating leukocyte subtype into the vessel wall [1]. Leukocyte adhesion to the vascular endothelium and subsequent transendothelial migration play essential roles in the pathogenesis of cardiovascular diseases such as atherosclerosis [2]. Several epidemiologic studies have reported that an increased leukocyte subtype count is a strong independent risk factor for cardiovascular (CV) events [3,4]. An increased leukocyte subtype count is an independent risk factor for the prevalence and progression of subclinical carotid atherosclerosis [5–11]. However, Kuo et al. [12] reported no association between total leukocyte count and IMT in asymptomatic subjects.

The different parameters used to assess vascular structure and function are related to the risk of cardiovascular morbidity and mortality. Similarly, ultrasound examination of the carotid arteries with the measurement of intima media thickness (IMT) or the presence of plaques has been shown to predict the occurrence of both stroke and myocardial infarction, independently of traditional CV risk factors [13,14]. The pulse wave velocity (PWV) has been associated with increased morbidity-mortality in both patients with cardiovascular disease and in healthy individuals [15,16]. The central augmentation index (CAIx) is an independent predictor of all-cause and CV mortality [17]. The Cardio-Ankle vascular index (CAVI) is also used to evaluate arterial stiffness and can be used to estimate the risk of atherosclerosis. This parameter is independent of arterial pressure at measuring time [18,19].

The association between leukocyte subtype counts and vascular structure and function in patients with intermediate risk has not been determined.

Moreover, there are differences in the measures used to assess the arterial stiffness according to gender [20], and discrepancies in the behavior of different measures are currently not completely resolved [21].

In patients with intermediate risk, it is important to analyze new cardiovascular risk factors and the association between them for personalized risk stratification [22].

Currently, many studies that analyze new techniques or biomarkers to improve risk estimation based on scales have focused on the population with intermediate cardiovascular risk [3,23,24].

Therefore, we investigated the relationship between leukocyte subtype counts and vascular structure and function based on carotid intima-media thickness, pulse wave velocity, central augmentation index and cardio-ankle vascular index in intermediate risk patients.

Methods

Study Design

This is a cross-sectional study to evaluate if ankle-brachial index (ABI), measures of arterial stiffness (CAVI), postprandial glucose, glycosylated hemoglobin, self-measured blood pressure and presence of comorbidity are independently associated to incidence of vascular events and whether they can improve the predictive capacity of current risk equations in the intermediate risk population. The second step will be a following of 5 and 10 years to evaluate cardiovascular morbidity and mortality. This study analyzed 500 subjects who were included at the baseline of MARK study (NCT01428934) [25].

Study Population

The population comprised individuals aged between 35 to 74 years who had intermediate cardiovascular risk, which was defined as coronary risk between 5% -15% at 10 years according to the Framingham adapted risk equation (REGICOR) [26] or vascular mortality risk between 2–5% at 10 years according to the SCORE equation [27]. The exclusion criteria were terminal illness, institutionalization at the appointment time, or a personal history of atherosclerotic disease. Sample selection was done with a random sample from the population aged 35 to 74 (both included) who had an intermediate cardiovascular risk. Recruitment and data collection for the study were carried out from July 2011 to June 2013. A sample-size calculation indicated that the 500 patients included in the study constituted a sufficient sample for detecting a correlation coefficient of 0.125 between leukocyte subtype counts and IMT in a two sided test, with a level of significance of 95% (alfa risk 0.05) and a power of 80% (beta risk 0.20). We have used the EPIDAT 4.0 software to perform this estimation. The study was approved by an independent ethics committee of Salamanca University Hospital (Spain), and all participants gave written informed consent according to the general recommendations of the Declaration of Helsinki [28].

Measurements

A detailed description has been published elsewhere regarding how the clinical data, anthropometric measurements, and analytical parameters were obtained [25].

Laboratory determinations

Venous blood sampling was performed between 08:00 and 09:00 after the individuals had fasted and abstained from smoking and the consumption of alcohol and caffeinated beverages for the previous 12 hours. Fasting plasma glucose, serum total cholesterol and high-density lipoprotein (HDL) cholesterol concentrations were measured using standard enzymatic automated methods. Low-density lipoprotein (LDL) cholesterol was estimated by the Friedewald equation when the direct parameter was not available. Non-HDL cholesterol was estimated by the equation (Non-HDL Cholesterol = Total Cholesterol- HDL Cholesterol). Blood samples were collected in the Alamedilla Health Center and analyzed at the hospital of Salamanca, which was approved by the external quality assurance programs of the Spanish Society of Clinical Chemistry and Molecular Pathology.

Office blood pressure

Office blood pressure (BP) was calculated as the average of the last two of three measurements of systolic blood pressure (SBP) and diastolic blood pressure (DBP) made with a validated sphygmomanometer (OMRON Model M10-IT). Measurements were made on the dominant

arm of participants in the seated position after at least 5 minutes of rest, with a cuff of appropriate size as determined by measurement of the upper arm circumference and following the recommendations of the European Society of Hypertension [29].

Central augmentation index (CAIx)

This parameter was estimated using an oscillometric Mobil-O-Graph (Stolberg, Germany) [30]. The device was validated according to British Hypertension Society [31] and European Society of Hypertension [32] recommendations. The measurements of central systolic blood pressure (cSBP) and peripheral systolic blood pressure (pSBP) were also taken on the dominant arm. Arm circumferences were measured and recorded to allow the correct choice of cuff size (two sizes available: 24–34 and 32–42 cm). With a conventional cuff, the determination of the cSBP is based on an oscillometric BP measurement and uses the pulse waves assessed at the brachial artery. After the estimation of peripheral BPs, the cuff instantly reinflates, and recordings for cSBP are carried out at diastolic pressure levels for 10 sec [33,34]. From the morphology of the aortic wave, CAIx was estimated using the following formula: increase in central pressure ×100/pulse pressure. The value was adjusted to a heart rate of 75 by the Mobil-O-Graph device.

Cardio Ankle Vascular Index (CAVI) and brachial ankle Pulse Wave Velocity (ba-PWV)

CAVI was measured using a Vasera VS-1500 device (Fukuda Denshi) and ba-PWV estimated by validate equation. The ba-PWV and CAVI were calculated to give a more accurate calculation of the degree of atherosclerosis. CAVI integrates cardiovascular elasticity derived from the aorta to the ankle pulse velocity through an oscillometric method. Takaki et al. [35] provided the evidence for the validity of CAVI by showing a positive correlation between the stiffness parameter β of the aorta and CAVI ($r = 0.67$, $p < 0.01$). The measurement of CAVI not affected by the increase in BP during measurement [18,19,36,37], CAVI values were automatically calculated by substituting the stiffness parameter β in the following equation to detect the vascular elasticity and the cardio-ankle PWV: Stiffness parameter $\beta = 2\rho \times 1/(P_s - P_d) \times \ln(P_s/P_d) \times PWV^2$, where ρ is the blood density, P_s and P_d are SBP and DBP in mmHg, and PWV is measured between the aortic valve and ankle. The average coefficient of variation of the CAVI is less than 5%, which is small enough for clinical use and confirms that CAVI has favorable reproducibility [18,38]. CAVI was measured at rest. CAVI was classified as normal (CAVI < 8), borderline (CAVI ≥ 8 and < 9) and abnormal (CAVI ≥ 9). Abnormal CAVI represents sub-clinical atherosclerosis. ba-PWV ≥ 17.5 m/sec was considered abnormal [39,40]. For the study, the higher obtained CAVI and ba-PWV were considered.

Assessment of vascular structure by carotid intima media thickness (IMT)

Carotid ultrasound was performed to assess carotid IMT by two investigators trained for this purpose before starting the study. The reliability of the recordings was evaluated before the study using the intra-class correlation coefficient, which showed values of 0.97 (95% CI: 0.94 to 0.99) for intra-observer agreement in repeated measurements of 20 subjects, and 0.90 (95% CI: 0.74 to 0.96) for inter-observer agreement. According to the Bland-Altman analysis, the mean difference for inter-observer agreement (95% limits of agreement) was 0.01 (-0.03 to 0.06). A Sonosite Micromax ultrasound device paired with a 5–10 MHz multi-frequency high-

resolution linear transducer with Sonocal software was used for performing automatic measurements of IMT in order to optimize reproducibility.

Measurements were made of the common carotid after the examination of a 10 mm longitudinal section at a distance of 1 cm from the bifurcation, performing measurements in the anterior and in the posterior wall, in the lateral, anterior and posterior projections, following an axis perpendicular to the artery to discriminate two lines: one for the intima-blood interface and the other for the media-adventitious interface. A total of 6 measurements of the right carotid were obtained, with another 6 measurements of the left carotid, using average values (average mean IMT and average maximum IMT) automatically calculated by the software [41]. The measurements were obtained with the subject lying down, with the head extended and slightly turned opposite to the examined carotid artery. Average mean IMT was considered abnormal if > 0.90 mm or if there were atherosclerotic plaques with a diameter of 1.5 mm or a focal increase of 0.5 mm or 50% of the adjacent IMT [42].

Anthropometric measurements

Body weight was determined on two occasions using a homologated electronic scale (Seca 770) following calibration (precision ± 0.1 kg), with the patient wearing light clothing and no shoes. Height in turn was measured with a portable system (Seca 222), recording the average of two readings. Body mass index (BMI) was calculated as weight (kg) divided by height squared (m^2). A value of > 30 kg/m^2 was taken to define obesity. The individuals performing the different tests were blinded to the clinical data of the patient. All assessments were made within a period of 10 days.

Statistical Analysis

Continuous variables were expressed as the mean \pm standard deviation for normally distributed continuous data, the median (interquartile range, IQR) for asymmetrically distributed continuous data, and the frequency distribution for categorical data. Statistical normality was tested using the Kolmogorov-Smirnov test. A Spearman's correlation was used to analyze the relationship between asymmetrically distributed continuous data. The difference of means between two categories of quantitative variables for asymmetrically distributed continuous data was analyzed using the Mann-Whitney U test. The difference of means between more than two categories of quantitative variables for symmetrically distributed continuous data was analyzed using the Anova test. We performed multiple linear regression analyses, one for each of the dependent variables, with IMT mean (x 100) and CAIx (x 10) to facilitate interpretation, ba-PWV and CAVI as dependent variables and total leukocyte count ($/mm^3$), monocyte count ($/mm^3$), neutrophil count ($/mm^3$) and lymphocyte count ($/mm^3$) as independent variables. We adjusted by age, systolic blood pressure, non-HDL cholesterol, fasting glucose, body mass index, number of cigarettes smoked, and antihypertensive, lipid-lowering and antidiabetic drugs. Comparisons between three or more groups were performed using ANOVA and differences between groups were assessed using DMS post hoc test. The data were analyzed using the Statistical Package for the Social Sciences version 20.0 (SPSS, Chicago, IL, USA). A value of $p < 0.05$ was considered statistically significant.

Results

The characteristics of the study subjects, global and by gender, are shown in [Table 1](#). The mean age was 60.3 ± 8.4 years, and 45.6% of the subjects were women. IMT and PWV had higher values in men, and CAIx was higher in women ($p < 0.01$). The total leukocyte, monocyte and neutrophil counts were higher in men.

Table 1. Baseline demographic and clinical characteristics of patients.

	Global n = 500	Women N = 228	Men N = 272	p-value
Age (years)	60.3±8.4	61.9±8.0	58.9±8.5	<0.001
Smoking n (%)	117 (23.4)	46 (39.3)	71(60.7)	0.138
Body mass index (kg/m ²)	27.8 (25.3–30.3)	27.5 (24.9–30.7)	27.9 (25.9–30.1)	0.254
Obesity n (%)	137 (27.4)	68 (49.6)	69 (50.4)	0.270
Office SBP (mmHg)	133.9±16.6	130.9±17.0	136.3±15.9	<0.001
Office DBP (mmHg)	81.3±10.7	79.3±11.2	82.9±10.1	<0.001
Heart Rate (beats/min)	69 (63–77)	70 (65–78)	67 (61–76)	0.004
Hypertension n (%)	401 (80.2)	175(43.6)	226 (56.4)	0.091
Antihypertensive Drugs n (%)	266 (53.2)	120 (45.1)	146 (54.9)	0.857
Fasting glucose (mg/dL)	89 (82–99)	88 (82–98)	90 (82–100)	0.226
Diabetes n (%)	138 (27.6)	51 (37.0)	87 (63.0)	0.021
Antidiabetic drugs n (%)	82 (16.4)	30 (36.6)	52 (63.4)	0.089
Total cholesterol (mg/dL)	216.2±38.5	221.1±36.8	212.1±39.3	0.009
LDL-cholesterol (mg/dL)	135.1±34.3	136.1±33.4	134.9±35.2	0.850
HDL-cholesterol (mg/dL)	53.5 (44.7–63.9)	59.9 (51.2–69.5)	49.0 (41.5–56.9)	<0.001
Non-HDL cholesterol (mg/dL)	80.4±14.0	84.3±13.3	77.1±13.8	<0.001
Dyslipidemia n (%)	418 (83.6)	197 (47.1)	221 (52.9)	0.146
Lipid lowering drugs n (%)	185 (37.0)	86 (37.7)	99 (36.4)	0.781
Total leukocyte count *10 ³ (/mm ³)	6.8 (5.7–8.1)	6.4 (5.4–7.6)	7.1 (6.1–8.3)	<0.001
Monocyte count *10 ³ (/mm ³)	0.5 (0.5–0.7)	0.5 (0.4–0.6)	0.6 (0.5–0.7)	<0.001
Neutrophil count *10 ³ (/mm ³)	3.5 (2.9–4.3)	3.3 (2.6–4.1)	3.7 (3.0–4.5)	<0.001
Lymphocyte count *10 ³ (/mm ³)	2.4 (2.0–2.9)	2.4 (2.0–2.8)	2.4 (1.9–3.0)	0.472
IMT mean (mm)	0.74±0.09	0.72±0.08	0.75±0.10	<0.001
TOD carotid n (%)	85 (17.1)	86 (46.5)	99 (53.5)	0.781
CAIx (%)	26.78±13.80	31.39±13.72	22.68±12.66	<0.001
ba-PWV (m/sec)	14.16 (12.69–16.19)	14.74 (12.83–16.46)	13.86(12.55–15.99)	0.009
ba-PWV ≥ 17.5 n (%)	65 (13.1)	35 (53.8)	30 (46.2)	0.142
CAVI	8.59±1.10	8.55±1.03	8.63±1.16	0.449
CAVI ≥ 9 n (%)	155 (31.3)	46 (39.3)	71 (60.7)	0.138

Values are means (standard deviations (SD)) for normally distributed continuous data and medians (interquartile range (IQR)) for asymmetrically distributed continuous data and number and proportions for categorical data. SBP: Systolic Blood Pressure. DBP: Diastolic Blood Pressure. HDL: High Density Lipoprotein. Non HDL-Colesterol = Total Cholesterol—LDL-Cholesterol. LDL: Low Density Lipoprotein. IMT: Intima Media Thickness of common carotid artery. TOD: Target Organ Damage. CAIx: Central Augmentation Index corrected for heart rate. ba-PWV: Brachial Ankle Pulse Wave Velocity. CAVI: Cardio-Ankle Vascular Index.

doi:10.1371/journal.pone.0119963.t001

[Table 2](#) shows the values of the total leukocyte, monocyte, neutrophil and lymphocyte counts according to the target organ damage of IMT, ba-PWV and CAVI. There was a greater total leukocyte count and in all series in subjects with pathologic IMT.

[Fig. 1](#) shows the values of the IMT by neutrophil count tertiles and the CAIx by monocyte count tertiles.

The total leukocyte, neutrophil and monocyte counts were positively correlated with mean average IMT ($p<0.01$) in men. The monocyte count was positively correlated with CAIx in women ($r = 0.212$; $p<0.01$) ([Table 3](#)). Simple linear regressions in [Fig. 2](#).

A multiple linear regression analysis by gender was carried out ([Table 4](#)). After adjusting for age, systolic blood pressure, non-HDL cholesterol, fasting glucose, body mass index, number of cigarettes smoked, antihypertensive drugs, lipid-lowering drugs and antidiabetic drugs, the

Table 2. Leukocyte subtypes counts according parameters of vascular structure and function.

TOD carotid	With TOD	Without TOD	p value
Total leukocyte count * 10^3 (/mm 3)	7.3 (6.2–8.6)	6.7 (5.6–7.9)	0.002
Monocyte count * 10^3 (/mm 3)	0.6 (0.5–0.7)	0.5 (0.4–0.6)	0.006
Neutrophil count * 10^3 (/mm 3)	3.7 (3.1–4.9)	3.5 (2.9–4.3)	0.005
Lymphocyte count (/mm 3)	2.6 (2.1–3.2)	2.4 (2.0–2.9)	0.059
ba-PWV \geq 17.5 m/sec			
Total leukocyte count * 10^3 (/mm 3)	6.9 (5.9–7.7)	6.7 (5.6–8.1)	0.711
Monocyte count * 10^3 (/mm 3)	0.5 (0.4–0.6)	0.5 (0.5–0.7)	0.700
Neutrophil count * 10^3 (/mm 3)	3.6 (2.9–4.4)	3.5 (2.9–4.3)	0.881
Lymphocyte count * 10^3 (/mm 3)	2.5 (2.1–3.2)	2.4 (2.0–2.9)	0.219
CAVI \geq 9			
Total leukocyte count * 10^3 (/mm 3)	6.7 (5.6–7.7)	6.8 (5.7–8.3)	0.291
Monocyte count * 10^3 (/mm 3)	0.5 (0.4–0.6)	0.5 (0.5–0.7)	0.922
Neutrophil count * 10^3 (/mm 3)	3.4 (2.9–4.2)	3.6 (2.9–4.4)	0.188
Lymphocyte count * 10^3 (/mm 3)	2.4 (2.0–2.9)	2.4 (2.0–2.9)	0.993

Values are medians (interquartile range (IQR)). TOD: Target Organ Damage. ba-PWV: Brachial Ankle Pulse Wave Velocity. CAVI: Cardio-Ankle Vascular Index. TOD carotid: Intima Media Thickness (IMT) > 0.90 mm or plaques with a diameter of 1.5 mm or a focal increase of 0.5 mm or 50% of the adjacent IMT

doi:10.1371/journal.pone.0119963.t002

mean IMT maintained a positive association with the neutrophil count ($\beta = 1.500$; $p = 0.007$) in men. CAIx maintained a positive association with the monocyte count ($\beta = 2.445$; $p = 0.022$) in women.

Discussion

The results of this study suggest that the relationship between subtype circulating leukocyte counts and vascular structure and function, although small, may be different by gender. The monocyte count was positively association with CAIx in women, and the neutrophil count was with IMT in men in a large sample of intermediate cardiovascular risk patients.

Several studies have demonstrated that neutrophils have an important role in the atherosclerosis development in the general population [10,43] and in men [7]. The monocytes are correlated with the IMT in healthy subjects [5,11] and in Japanese subjects with type 2 diabetes [4].

The results of this study are consistent with those previously disclosed and add new information. According to our results, this correlation is observed only in men, and suggest that the role of neutrophils may be more important than monocytes when IMT is increased in men with mid-level cardiovascular risk. Our results support the paper published by Li et al. [44], who found a relation only in men between a high number of leucocytes and a high incidence rate of cardio vascular diseases, but not with stroke. In summary, the leukocytes seem to have an independent role in early arterial damage and they may reflect subclinical disease in men.

As far as we know, this is the first study that describes a positive association between monocytes and CAIx in women. The assessment of CAIx is a simple approach to quantify the role of wave reflection in determining an elevation of central blood pressure values [16,45]. This relationship between CAIx and monocytes in women opens new lines of research.

The differences found between the two genders may be due to several factors. First, IMT is higher in men, while CAIx is higher in women. The influence of sex steroids on vascular function has been demonstrated in association with hormonal changes throughout the lifespan




Fig 1. Mean IMT values according to the neutrophil count tertiles in men; Neutrophil count tertiles: T1 < 3000/mm³; T2 3000 to 4000/mm³; T3 > 4000/mm³. CAIx in women values according to the monocyte count tertiles. Monocyte/count tertiles: 1° Tertile < 500 /mm³, 2° Tertile 500 to 600/mm³, 3° Tertile > 600 /mm³. ANOVA Test: Carotid intima media thickness by neutrophil count tertiles in men: p = 0.035; CAIx by monocyte count tertiles in women: p = 0.042.

doi:10.1371/journal.pone.0119963.g001

(puberty in both genders and menopause in females) [20,46]. Furthermore, anthropometric factors such as distribution of body fat [47], height, and aortic length are different between

Table 3. Bivariate correlations of vascular structure and function indices with leukocyte subtype counts.

	Total leukocyte count (/mm ³)	Monocyte count (/mm ³)	Neutrophil count (/mm ³)	Lymphocyte count (/mm ³)
IMT mean:				
Global	0.104*	0.128**	0.103*	0.089*
Men	0.167**	0.152*	0.170**	0.093
Women	-0.010	0.015	-0.019	0.086
CAIx				
Global	0.027	0.017	-0.007	0.076
Men	0.062	0.030	0.037	0.070
Women	0.116	0.212**	0.048	0.127
ba-PWV:				
Global	0.039	-0.019	0.045	0.069
Men	0.081	0.067	0.078	0.086
Women	0.035	-0.053	0.055	0.056
CAVI				
Global	-0.049	-0.012	-0.058	0.014
Men	-0.056	0.012	-0.053	-0.019
Women	-0.041	-0.050	-0.067	0.052

IMT: Intima Media Thickness of common carotid artery. CAIx: Central Augmentation Index corrected for heart rate. ba-PWV: Brachial Ankle Pulse Wave Velocity. CAVI: Cardio-Ankle Vascular Index.

p-values by Spearman correlation.

* p<0.05

** p<0.01.

doi:10.1371/journal.pone.0119963.t003

men and women [20,48,49]. Finally, the role of inflammatory components in the development of atherosclerosis is likely to differ according to gender. Thus, in a study conducted by our group in hypertensive patients, we found that hs-CRP has a positive association with IMT in men and a negative association with CAIx in women [50]. This result suggests that gender differences in cardiovascular outcomes may be mediated in part via gender differences in artery wall properties throughout life and that further understanding may lead to improved strategies for the prevention of cardiovascular diseases in both women and men.

The relation between the ba-PWV with the leukocyte subtype counts is not clear. In the same line as our results, Kavousiet al. [4] found that ba-PWV was not correlated with monocyte count, whereas Phillips et al. [11] found that leukocyte count ($\beta = 0.07$; $p < 0.001$) and granulocyte number ($\beta = 0.07$; $p < 0.001$) were significantly positively related to ba-PWV in a Chinese population, but not lymphocyte number. Bearing in mind these opposite results, it is considered as necessary to carry out more prospective studies in order to elucidate the association between ba-PWV and leukocyte subtype counts and the differences that could be observed in each gender.

The only study that has been found that analyzed the relationship between subtype circulating leukocyte counts with CAVI in 3738 Japanese people in the general population showed that the leukocyte counts weakly correlated with CAVI in men ($\beta = 0.61$; $p = 0.043$), but not in women ($\beta = 0.35$; $p = 0.17$) [51]. Our study has not shown any correlation between the leukocyte counts or their subtypes with the CAVI in neither men nor women.

Limitations

This study has several limitations what should be considered in the interpretation of our results. Because this was a cross-sectional, observational study, which prevented us from




Fig 2. IMT regression line with Neutrophil count: Women: ($r = -0.019, p = 0.781$); Men: ($r = 0.170, p = 0.005$); CAIx regression line with monocyte count: Women: ($r = 0.212, p = 0.001$); Men: ($r = 0.030, p = 0.629$).

doi:10.1371/journal.pone.0119963.g002

Table 4. Multiple Regression Analysis with vascular structure and function parameters as dependent variable and leukocyte subtype counts as independent variable.

Dependent variable:	Men			Women		
	β	CI 95%	p value	β	CI 95%	p value
IMT mean						
Total leukocyte count (/mm ³)	0.721	0.002 to 1.439	0.049	0.234	-0.590 to 1.058	0.574
Monocyte count (/mm ³)	2.281	-4.713 to 9.274	0.521	7.007	-3.887 to 17.901	0.204
Neutrophil count (/mm ³)	1.500	0.409 to 2.592	0.007	0.247	-1.049 to 1.543	0.705
Lymphocyte count (/mm ³)	0.225	-1.369 to 1.818	0.781	0.651	-1.942 to 3.243	0.619
CAIx						
Total leukocyte count (/mm ³)	0.081	-0.023 to 0.185	0.128	0.028	-0.133 to 0.190	0.726
Monocyte count (/mm ³)	0.225	-0.785 to 1.237	0.660	2.445	0.359 to 4.530	0.022
Neutrophil count (/mm ³)	0.076	-0.085 to 0.237	0.355	-0.014	-0.268 to 0.240	0.912
Lymphocyte count (/mm ³)	0.223	-0.004 to 0.450	0.055	-0.029	-0.539 to 0.479	0.908
ba-PWV						
Total leukocyte count (/mm ³)	0.130	-0.082 to 0.342	0.229	-0.016	-0.213 to 0.180	0.869
Monocyte count (/mm ³)	0.955	-1.091 to 3.002	0.358	-1.346	-3.950 to 1.257	0.306
Neutrophil count (/mm ³)	0.169	-0.156 to 0.494	0.308	-0.002	-0.310 to 0.307	0.991
Lymphocyte count (/mm ³)	0.288	-0.176 to 0.753	0.222	0.352	-0.261 to 0.964	0.257
CAVI						
Total leukocyte count (/mm ³)	0.025	-0.050 to 0.101	0.508	-0.047	-0.143 to 0.49	0.335
Monocyte count (/mm ³)	0.036	-0.693 to 0.765	0.922	-1.009	-2.276 to 0.257	0.117
Neutrophil count (/mm ³)	0.019	-0.097 to 0.135	0.745	-0.060	-0.211 to 0.091	0.428
Lymphocyte count (/mm ³)	0.103	-0.063 to 0.268	0.222	0.143	-0.159 to 0.444	0.349

Dependent variables: IMT mean: Intima Media thickness of common carotid artery. CAIx: Central Augmentation Index corrected for heart rate. ba-PWV: Brachial ankle Pulse Wave Velocity. CAVI: Cardio-Ankle Vascular Index. **Independent variables:** Total leukocyte count. Monocyte count. Neutrophil count. Lymphocyte count. **Adjusted by:** Age, systolic blood pressure, Non-high density lipoprotein, fasting glucose, body mass index, number of cigarettes, antihypertensive drugs, lipid lowering drugs and antidiabetic drugs.

doi:10.1371/journal.pone.0119963.t004

establishing a temporal relationship between the circulating leukocyte subtype counts and the different parameters used to assess vascular function and structure and not allow us to establish a causal relationship. Some subjects were on treatment with antihypertensive or lipid lowering drugs, which might have affected the arterial stiffness measures. However, we adjusted for anti-hypertensive, lipid lowering and antidiabetic drugs. Finally, the correlations were small and need to be confirmed in longitudinal studies the clinical importance of them. However, this is the first study to examine the relationship between the circulating leukocyte subtype counts and different vascular structure and function parameters in intermediate-risk patients.

Conclusions

The results of this study suggest that the relationship between subtype circulating leukocyte counts and vascular structure and function, although small, may be different by gender. In men, the neutrophil count was positively correlated with IMT and in women, the monocyte count with CAIx, in a large sample of intermediate-risk patients. These associations were maintained after adjusting for age and other confounders.

Further research involving prospective studies are required to define the role of these leukocyte subtype counts in the vascular structure and function. In addition, possible gender-specific roles of leukocyte subtype counts remain to be elucidated.

Supporting Information

S1 Data.

(SAV)

Acknowledgments

We are grateful to all professionals participating in the MARK study.

Lead author for this group: Rafel Ramos: Research Unit, Primary Health Care, Girona. Jordi Gol Institute for Primary Care Research (IDIAP Jordi Gol), Girona, Spain. E-mail: raramos.girona.ics@gencat.net. Coordinating Center: Rafel Ramos, Ruth Martí, Dídac Parramon, Anna Ponjoan, Miquel Quesada, Maria Garcia-Gil, Martina Sidera and Lourdes Camós. Research Unit, Primary Health Care. Jordi Gol Institute for Primary Care Research (IDIAP Jordi Gol). C/Maluquer Salvador, 11. 17002-Girona. Girona, Spain. Fernando Montesinos, Ignacio Montoya, Carlos López, Anna Agell, Núria Pagès of the Primary Care Services, Girona. Catalan Institute of Health (ICS), Catalonia, Spain. Irina Gil, Anna Maria-Castro of the Primary Care Services, Girona. Institut d'Assistència Sanitaria (IAS), Catalonia, Spain. Fernando Rigo, Guillermo Frontera, Antònia Rotger, Natalia Feuerbach, Susana Pons, Natividad Garcia, John Guillamet, Micaela Llull and Mercedes Gutierrez of the Health Center Primary Care San Agustín. Ibsalut Balears, Spain. Cristina Agudo-Conde, Leticia Gomez-Sánchez, Carmen Castaño-Sánchez, Carmela Rodriguez-Martín, Benigna Sanchez-Salgado, Angela de Cabo-Laso, Marta Gomez-Sánchez, Emiliano Rodriguez-Sánchez, Jose Angel Maderuelo-Fernandez, Emilio Ramos-Delgado, Carmen Patino-Alonso, Jose I Recio-Rodriguez, Manuel A Gomez-Marcos and Luis Garcia-Ortiz. Primary Care Research Unit of the Alamedilla, Salamanca, Spain. Castilla and León Health Service-SACYL

Author Contributions

Conceived and designed the experiments: MAGM RR LGO RM. Performed the experiments: JIRR LGS CAC. Analyzed the data: MCPA MAGM LGO. Contributed reagents/materials/analysis tools: JIRR CAC FR. Wrote the paper: LGS LGO JIRR RR RM MAGM.

References

1. Tuttolomondo A, Di Raimondo D, Pecoraro R, Arnao V, Pinto A, Licata G: Atherosclerosis as an inflammatory disease. *Curr Pharm Des* 2012; 18:4266–4288. PMID: [22390643](https://pubmed.ncbi.nlm.nih.gov/22390643/)
2. Zhang J, Alcaide P, Liu L, Sun J, He A, Luscinskas FW, et al.: Regulation of endothelial cell adhesion molecule expression by mast cells, macrophages, and neutrophils. *PLoS One* 2011; 6:e14525. doi: [10.1371/journal.pone.0014525](https://doi.org/10.1371/journal.pone.0014525) PMID: [21264293](https://pubmed.ncbi.nlm.nih.gov/21264293/)
3. Kavousi M, Elias-Smale S, Rutten JH, Leening MJ, Vliegenthart R, Verwoert GC, et al.: Evaluation of newer risk markers for coronary heart disease risk classification: a cohort study. *Ann Intern Med* 2012; 156:438–444. doi: [10.7326/0003-4819-156-6-201203200-00006](https://doi.org/10.7326/0003-4819-156-6-201203200-00006) PMID: [22431676](https://pubmed.ncbi.nlm.nih.gov/22431676/)
4. Matsumura T, Taketa K, Motoshima H, Senokuchi T, Ishii N, Kinoshita H, et al.: Association between circulating leukocyte subtype counts and carotid intima-media thickness in Japanese subjects with type 2 diabetes. *Cardiovasc Diabetol* 2013; 12:177. doi: [10.1186/1475-2840-12-177](https://doi.org/10.1186/1475-2840-12-177) PMID: [24373412](https://pubmed.ncbi.nlm.nih.gov/24373412/)
5. Chapman CM, Beilby JP, McQuillan BM, Thompson PL, Hung J: Monocyte count, but not C-reactive protein or interleukin-6, is an independent risk marker for subclinical carotid atherosclerosis. *Stroke* 2004; 35:1619–1624. PMID: [15155967](https://pubmed.ncbi.nlm.nih.gov/15155967/)
6. Johnsen SH, Fosse E, Joakimsen O, Mathiesen EB, Stensland-Bugge E, Njolstad I, et al.: Monocyte count is a predictor of novel plaque formation: a 7-year follow-up study of 2610 persons without carotid plaque at baseline the Tromso Study. *Stroke* 2005; 36:715–719. PMID: [15746459](https://pubmed.ncbi.nlm.nih.gov/15746459/)
7. Loimaa A, Rontu R, Vuori I, Mercuri M, Lehtimaki T, Nenonen A, et al.: Blood leukocyte count is a risk factor for intima-media thickening and subclinical carotid atherosclerosis in middle-aged men. *Atherosclerosis* 2006; 188:363–369. PMID: [16378612](https://pubmed.ncbi.nlm.nih.gov/16378612/)

8. Nozawa N, Hibi K, Endo M, Sugano T, Ebina T, Kosuge M, et al.: Association between circulating monocytes and coronary plaque progression in patients with acute myocardial infarction. *Circ J* 2010; 74:1384–1391. PMID: [20467155](#)
9. Lau KK, Wong YK, Chan YH, Yiu KH, Teo KC, Li LS, et al.: Prognostic implications of surrogate markers of atherosclerosis in low to intermediate risk patients with type 2 diabetes. *Cardiovasc Diabetol* 2012; 11:101. doi: [10.1186/1475-2840-11-101](#) PMID: [22900680](#)
10. Ortega E, Gilabert R, Nunez I, Cofan M, Sala-Vila A, de Groot E, et al.: White blood cell count is associated with carotid and femoral atherosclerosis. *Atherosclerosis* 2012; 221:275–281. doi: [10.1016/j.atherosclerosis.2011.12.038](#) PMID: [22244768](#)
11. Phillips AC, Jiang CQ, Thomas GN, Lin JM, Yue XJ, Cheng KK, et al.: White blood cell subsets are associated with carotid intima-media thickness and pulse wave velocity in an older Chinese population: the Guangzhou Biobank Cohort Study. *J Hum Hypertens* 2012; 26:485–492. doi: [10.1038/jhh.2011.58](#) PMID: [21654852](#)
12. Kuo WK, Lee SY, Ma SM, Ling TA, Wu CC: Correlation of hematologic factors to carotid intima-media thickness in men and women: a study of 2767 asymptomatic subjects of Taiwan. *Acta Neurol Taiwan* 2012; 21:158–164. PMID: [23329546](#)
13. O'Leary DH, Polak JF, Kronmal RA, Manolio TA, Burke GL, Wolfson SK Jr.: Carotid-artery intima and media thickness as a risk factor for myocardial infarction and stroke in older adults. *Cardiovascular Health Study Collaborative Research Group*. *N Engl J Med* 1999; 340:14–22. PMID: [9878640](#)
14. Nambi V, Chambliss L, Folsom AR, He M, Hu Y, Mosley T, et al.: Carotid intima-media thickness and presence or absence of plaque improves prediction of coronary heart disease risk: the ARIC (Atherosclerosis Risk In Communities) study. *J Am Coll Cardiol* 2010; 55:1600–1607. doi: [10.1016/j.jacc.2009.11.075](#) PMID: [20378078](#)
15. Mattace-Raso FU, van der Cammen TJ, Hofman A, van Poppel NM, Bos ML, Schalekamp MA, et al.: Arterial stiffness and risk of coronary heart disease and stroke: the Rotterdam Study. *Circulation* 2006; 113:657–663. PMID: [16461838](#)
16. Laurent S, Cockcroft J, Van Bortel L, Boutouyrie P, Giannattasio C, Hayoz D, et al.: Expert consensus document on arterial stiffness: methodological issues and clinical applications. *Eur Heart J* 2006; 27:2588–2605. PMID: [17000623](#)
17. Vlachopoulos C, Aznaouridis K, Stefanadis C: Prediction of cardiovascular events and all-cause mortality with arterial stiffness: a systematic review and meta-analysis. *J Am Coll Cardiol* 2010; 55:1318–1327. doi: [10.1016/j.jacc.2009.10.061](#) PMID: [20338492](#)
18. Shirai K, Utino J, Otsuka K, Takata M: A novel blood pressure-independent arterial wall stiffness parameter; cardio-ankle vascular index (CAVI). *J Atheroscler Thromb* 2006; 13:101–107. PMID: [16733298](#)
19. Shirai K, Hiruta N, Song M, Kurosu T, Suzuki J, Tomaru T, et al.: Cardio-ankle vascular index (CAVI) as a novel indicator of arterial stiffness: theory, evidence and perspectives. *J Atheroscler Thromb* 2011; 18:924–938. PMID: [21628839](#)
20. Rossi P, Frances Y, Kingwell BA, Ahimastos AA: Gender differences in artery wall biomechanical properties throughout life. *J Hypertens* 2011; 29:1023–1033. doi: [10.1097/HJH.0b013e328344da5e](#) PMID: [21346620](#)
21. Vermeersch SJ, Rietzschel ER, De Buyzere ML, De Bacquer D, De Backer G, Van Bortel LM, et al.: Age and gender related patterns in carotid-femoral PWV and carotid and femoral stiffness in a large healthy, middle-aged population. *J Hypertens* 2008; 26:1411–1419. doi: [10.1097/HJH.0b013e3282ffac00](#) PMID: [18551018](#)
22. Zhu J, Chen T, Yang L, Li Z, Wong MM, Zheng X, et al.: Regulation of microRNA-155 in atherosclerotic inflammatory responses by targeting MAP3K10. *PLoS One* 2012; 7:e46551. doi: [10.1371/journal.pone.0046551](#) PMID: [23189122](#)
23. Greenland P, Alpert JS, Beller GA, Benjamin EJ, Budoff MJ, Fayad ZA, et al.: 2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Circulation* 2010; 122:e584–636. doi: [10.1161/CIR.0b013e3182051b4c](#) PMID: [21098428](#)
24. Yeboah J, McClelland RL, Polonsky TS, Burke GL, Sibley CT, O'Leary D, et al.: Comparison of novel risk markers for improvement in cardiovascular risk assessment in intermediate-risk individuals. *JAMA* 2012; 308:788–795. doi: [10.1001/jama.2012.9624](#) PMID: [22910756](#)
25. Martí R, Parramon D, García-Ortiz L, Rigo F, Gómez-Marcos MA, Sempere I, et al.: Improving interMediate risk management. MARK study. *BMC Cardiovasc Disord* 2011; 11:61. doi: [10.1186/1471-2261-11-61](#) PMID: [21992621](#)
26. Marrugat J, D'Agostino R, Sullivan L, Elosua R, Wilson P, Ordovas J, et al.: An adaptation of the Framingham coronary heart disease risk function to European Mediterranean areas. *J Epidemiol Community Health* 2003; 57:634–638. PMID: [12883073](#)

27. Conroy RM, Pyorala K, Fitzgerald AP, Sans S, Menotti A, De Backer G, et al.: Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project. *Eur Heart J* 2003; 24:987–1003. PMID: [12788299](#)
28. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA* 2013; 310:2191–2194. doi: [10.1001/jama.2013.281053](#) PMID: [24141714](#)
29. O'Brien E, Asmar R, Beilin L, Imai Y, Mancia G, Mengden T, et al.: Practice guidelines of the European Society of Hypertension for clinic, ambulatory and self blood pressure measurement. *J Hypertens* 2005; 23:697–701. PMID: [15775768](#)
30. Wasertheurer S, Kropf J, Weber T, van der Giet M, Baulmann J, Ammer M, et al.: A new oscillometric method for pulse wave analysis: comparison with a common tonometric method. *J Hum Hypertens* 2010; 24:498–504. doi: [10.1038/jhh.2010.27](#) PMID: [20237499](#)
31. Wei W, Tolle M, Zidek W, van der Giet M: Validation of the mobil-O-Graph: 24 h-blood pressure measurement device. *Blood Press Monit* 2010; 15:225–228. doi: [10.1097/MBP.0b013e328338892f](#) PMID: [20216407](#)
32. Franssen PM, Imholz BP: Evaluation of the Mobil-O-Graph new generation ABPM device using the ESH criteria. *Blood Press Monit* 2010; 15:229–231. PMID: [20658764](#)
33. Protopero AD, Argyris A, Nasothimiou E, Vrachatis D, Papaioannou TG, Tzamouranis D, et al.: Feasibility and reproducibility of noninvasive 24-h ambulatory aortic blood pressure monitoring with a brachial cuff-based oscillometric device. *Am J Hypertens* 2012; 25:876–882. doi: [10.1038/ajh.2012.63](#) PMID: [22673021](#)
34. Weiss W, Gohlisch C, Harsch-Gladisch C, Tolle M, Zidek W, van der Giet M: Oscillometric estimation of central blood pressure: validation of the Mobil-O-Graph in comparison with the SphygmoCor device. *Blood Press Monit* 2012; 17:128–131. doi: [10.1097/MBP.0b013e328353ff63](#) PMID: [22561735](#)
35. Takaki A, Ogawa H, Wakeyama T, Iwami T, Kimura M, Hadano Y, et al.: Cardio-ankle vascular index is a new noninvasive parameter of arterial stiffness. *Circ J* 2007; 71:1710–1714. PMID: [17965489](#)
36. Shirai K, Song M, Suzuki J, Kurosu T, Oyama T, Nagayama D, et al.: Contradictory effects of beta1- and alpha1-adrenergic receptor blockers on cardio-ankle vascular stiffness index (CAVI)—CAVI independent of blood pressure. *J Atheroscler Thromb* 2011; 18:49–55. PMID: [21071883](#)
37. Takaki A, Ogawa H, Wakeyama T, Iwami T, Kimura M, Hadano Y, et al.: Cardio-ankle vascular index is superior to brachial-ankle pulse wave velocity as an index of arterial stiffness. *Hypertens Res* 2008; 31:1347–1355. doi: [10.1291/hypres.31.1347](#) PMID: [18957805](#)
38. Kubozono T, Miyata M, Ueyama K, Nagaki A, Otsuji Y, Kusano K, et al.: Clinical significance and reproducibility of new arterial distensibility index. *Circ J* 2007; 71:89–94. PMID: [17186984](#)
39. Korkmaz L, Erkan H, Korkmaz AA, Acar Z, Agac MT, Bektas H, et al.: Relationship of aortic knob width with cardio-ankle vascular stiffness index and its value in diagnosis of subclinical atherosclerosis in hypertensive patients: a study on diagnostic accuracy. *Anadolu Kardiyol Derg* 2012; 12:102–106. doi: [10.5152/akd.2012.034](#) PMID: [22281788](#)
40. Kawai T, Ohishi M, Onishi M, Ito N, Takeya Y, Maekawa Y, et al.: Cut-off value of brachial-ankle pulse wave velocity to predict cardiovascular disease in hypertensive patients: a cohort study. *J Atheroscler Thromb* 2013; 20:391–400. PMID: [23268984](#)
41. Gomez-Marcos MA, Recio-Rodriguez JL, Patino-Alonso MC, Agudo-Conde C, Gomez-Sanchez L, Gomez-Sanchez M, et al.: Protocol for measuring carotid intima-media thickness that best correlates with cardiovascular risk and target organ damage. *Am J Hypertens* 2012; 25:955–961. doi: [10.1038/ajh.2012.72](#) PMID: [22717546](#)
42. Mancia G, Fagard R, Narkiewicz K, Redon J, Zanchetti A, Bohm M, et al.: 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens* 2013; 31:1281–1357. doi: [10.1097/01.hjh.0000431740.32696.cc](#) PMID: [23817082](#)
43. Soehnlein O: Multiple roles for neutrophils in atherosclerosis. *Circ Res* 2012; 110:875–888. doi: [10.1161/CIRCRESAHA.111.257535](#) PMID: [22427325](#)
44. Li C, Engstrom G, Hedblad B: Leukocyte count is associated with incidence of coronary events, but not with stroke: a prospective cohort study. *Atherosclerosis* 2010; 209:545–550. doi: [10.1016/j.atherosclerosis.2009.09.029](#) PMID: [19833340](#)
45. McEnery CM, Yasmin, Maki-Petaja KM, McDonnell BJ, Munnery M, Hickson SS, et al.: The impact of cardiovascular risk factors on aortic stiffness and wave reflections depends on age: the Anglo-Cardiff Collaborative Trial (ACCT III). *Hypertension* 2010; 56:591–597. doi: [10.1161/HYPERTENSIONAHA.110.156950](#) PMID: [20696989](#)

46. Marlatt KL, Kelly AS, Steinberger J, Dengel DR: The influence of gender on carotid artery compliance and distensibility in children and adults. *J Clin Ultrasound* 2013; 41:340–346. doi: [10.1002/jcu.22015](https://doi.org/10.1002/jcu.22015) PMID: [23233368](#)
47. Anoop S, Misra A, Bhardwaj S, Gulati S: High body fat and low muscle mass are associated with increased arterial stiffness in Asian Indians in North India. *J Diabetes Complications* 2014.
48. McEnery CM, Yasmin, Hall IR, Qasem A, Wilkinson IB, Cockcroft JR: Normal vascular aging: differential effects on wave reflection and aortic pulse wave velocity: the Anglo-Cardiff Collaborative Trial (ACCT). *J Am Coll Cardiol* 2005; 46:1753–1760. PMID: [16256881](#)
49. Yasmin, Brown MJ: Similarities and differences between augmentation index and pulse wave velocity in the assessment of arterial stiffness. *QJM* 1999; 92:595–600. PMID: [10627881](#)
50. Gomez-Marcos MA, Recio-Rodriguez JI, Patino-Alonso MC, Agudo-Conde C, Gomez-Sanchez L, Rodriguez-Sanchez E, et al.: Relationships between high-sensitive C-reactive protein and markers of arterial stiffness in hypertensive patients. Differences by sex. *BMC Cardiovasc Disord* 2012; 12:37. doi: [10.1186/1471-2261-12-37](#) PMID: [22676422](#)
51. Sekitani Y, Hayashida N, Kadota K, Yamasaki H, Abiru N, Nakazato M, et al.: White blood cell count and cardiovascular biomarkers of atherosclerosis. *Biomarkers* 2010; 15:454–460. doi: [10.3109/1354750X.2010.486870](#) PMID: [20507253](#)