

CORRECTION

Correction: Exploratory Analysis on the Use of Statins with or without n-3 PUFA and Major Events in Patients Discharged for Acute Myocardial Infarction: An Observational Retrospective Study

The *PLOS ONE* Staff

There is an error in [Table 4](#). The Risk ratios for the last row, “Death or stroke” should read as follows: 0.65 (0.58 to 0.73).

Please find the complete, correct [Table 4](#) here:



OPEN ACCESS

Citation: The *PLOS ONE* Staff (2015) Correction: Exploratory Analysis on the Use of Statins with or without n-3 PUFA and Major Events in Patients Discharged for Acute Myocardial Infarction: An Observational Retrospective Study. *PLoS ONE* 10(4): e0126303. doi:10.1371/journal.pone.0126303

Published: April 17, 2015

Copyright: © 2015 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 4. Baseline characteristics and outcomes in paired-matched population.

Variable	Statins	Statins plus n-3 PUFA	P value*	
N	3,962	3,962	-	
Age, mean (SD)	65.1 (10.2)	65.2 (9.9)	0.29	
Male sex, n (%)	2,909 (73.4)	2,918 (73.6)	0.828	
Diabetes, n (%)	1,094 (27.6)	1,088 (27.5)	0.899	
Hypertension, n (%)	2,643 (66.7)	2,673 (67.5)	0.465	
Previous MI, n (%)	194 (4.9)	184 (4.6)	0.637	
Previous CHF, n (%)	476 (12.0)	476 (12.0)	1.00	
Previous Stroke, n (%)	69 (1.7)	63 (1.6)	0.656	
Previous AF, n (%)	209 (5.3)	219 (5.5)	0.643	
Peripheral vascular disease, n (%)	256 (6.5)	250 (6.3)	0.821	
Previous CABG, n (%)	36 (0.9)	37 (0.9)	1.00	
Previous PTCA, n (%)	174 (4.4)	167 (4.2)	0.742	
COPD, n (%)	402 (10.1)	395 (10.0)	0.815	
Malignancy, n (%)	88 (2.2)	93 (2.3)	0.764	
Depression, n (%)	106 (2.7)	104 (2.6)	0.944	
<i>In-hospital complications and procedures</i>				
Heart failure, n (%)	358 (9.0)	354 (8.9)	0.906	
AF, n (%)	164 (4.1)	173 (4.4)	0.650	
Coronary angiography, n (%)	1,852 (46.7)	1,799 (45.4)	0.240	
PTCA, n (%)	834 (21.0)	844 (21.3)	0.805	
CABG, n (%)	67 (1.7)	77 (1.9)	0.447	
<i>Treatments after discharge</i>				
Beta-blockers, n (%)	3,305 (83.4)	3,299 (83.3)	0.869	
ACEi / ARB, n (%)	3,539 (89.3)	3,554 (89.7)	0.600	
Clopidogrel, n (%)	1,714 (43.3)	1,667 (42.1)	0.294	
Aspirin, n (%)	3,681 (92.9)	3,666 (92.5)	0.534	
Outcomes				
	Statins	Statins plus n-3 PUFA	Risk ratios (95% CIs)	P value
All cause death	539 (13.6%)	340 (8.6%)	0.63 (0.56 to 0.72)	<0.001
Death or myocardial infarction	848 (21.4)	804 (20.3)	0.95 (0.87 to 1.03)	0.234
Death or atrial fibrillation	805 (20.3)	660 (16.7)	0.82 (0.75 to 0.90)	<0.001
Death or CHF	792 (20.0)	684 (17.3)	0.86 (0.79 to 0.95)	0.002
Death or stroke	662 (16.7)	456 (11.5)	0.65 (0.58 to 0.73)	<0.001

*All p values are for McNemar's test

doi:10.1371/journal.pone.0126303.t001

Reference

1. Macchia A, Romero M, D'Ettoire A, Tognoni G, Mariani J (2013) Exploratory Analysis on the Use of Statins with or without n-3 PUFA and Major Events in Patients Discharged for Acute Myocardial Infarction: An Observational Retrospective Study. PLoS ONE 8(5): e62772. doi: [10.1371/journal.pone.0062772](https://doi.org/10.1371/journal.pone.0062772) PMID: [23671633](https://pubmed.ncbi.nlm.nih.gov/23671633/)