

Gastrointestinal medication group	AN	BN	EDNOS-P	BED	No ED
	(n=54)	(n=585)	(n=61)	(n=3104)	(n=58215)
	<i>n (%)</i>	<i>n (%)</i>	<i>n (%)</i>	<i>n (%)</i>	<i>n (%)</i>
Antacids					
Before pregnancy	-	2 (0.3)	-	5 (0.2)	81 (0.1)
First trimester	-	7 (1.2)	1 (1.6)	40 (1.3)	551 (0.9)
Second trimester	2 (3.7)	37 (6.3)	7 (11.5)	266 (8.6)*	3536 (6.1)
Third trimester	6 (11.1)	43 (7.4)	9 (14.8)	284 (9.1)*	3923 (6.7)
Any time during pregnancy	6 (11.1)	49 (8.4)	10 (16.4)	348 (11.2)*	4805 (8.3)
0-3 months postpartum	-	5 (0.9)‡	-	14 (0.5)	135 (0.2)
4-6 months postpartum	-	6 (1.0)‡	-	11 (0.4)	124 (0.2)
Before, during and after pregnancy	-	-	-	-	6 (0.01)
Drugs for peptic ulcer and GERD					
Before pregnancy	-	5 (0.9)	-	29 (0.9)	453 (0.8)
First trimester	1 (1.9)	7 (1.2)	1 (1.6)	57 (1.8)*	654 (1.1)
Second trimester	6 (11.1)	67 (11.5)	6 (9.8)	344 (11.1)*	5258 (9.0)
Third trimester	6 (11.1)	74 (12.6)	6 (9.8)	394 (12.7)*	6315 (10.8)
Any time during pregnancy	7 (13.0)	92 (15.7)	8 (13.1)	473 (15.2)*	7503 (12.9)
0-3 months postpartum	-	2 (0.3)	-	19 (0.6)‡	184 (0.3)
4-6 months postpartum	-	-	-	19 (0.6)‡	173 (0.3)
Before, during and after pregnancy	-	-	-	2 (0.1)	27 (0.05)
Laxatives					
Before pregnancy	-	5 (0.9)*	1 (1.6)	1 (0.03)	76 (0.1)
First trimester	2 (3.7)	21 (3.6)*	-	67 (2.2)	936 (1.6)
Second trimester	5 (9.3)	42 (7.2)*	3 (4.9)	154 (5.0)‡	2247 (3.9)

Third trimester	4 (7.4)	35 (6.0)[*]	5 (8.2)	107 (3.4)	1738 (3.0)
Any time during pregnancy	6 (11.1)	61 (10.4)[*]	6 (9.8)	215 (6.9)[‡]	3279 (5.6)
0-3 months postpartum	5 (9.3)	31 (5.3)[‡]	1 (1.6)	112 (3.6)	1945 (3.3)
4-6 months postpartum	3 (5.6)[‡]	12 (2.1)[*]	-	30 (1.0)	406 (0.7)
Before, during and after pregnancy	-	-	-	-	12 (0.02)
