

Table S2: Primers used for qRT-PCR.

Target gene	Forward primer	Reverse primer	Size of amplicon (bp)
<i>cpa</i>	AATATGCTGAAGGTGACTACTC	TATGTACCATCCGACAACTGC	138
<i>emm53</i>	AATAGAGCAGACGACGCTAG	TTCCCGTAAATCAGTCTTAAGC	132
<i>fctA</i>	TTGAACCACTACTACTGCAAG	AGCAGACTCTTTATCTGTATTG	121
<i>fctB</i>	AGACAGCACTGTTCAAACCTAGC	TCGTCTATTGTTTTTCATCGCATC	113
<i>hsp33</i>	ATGGCACAGGTAACCCCTTACACCTC	ACATTAAGTCCGATTGCAGATGGTG	117
<i>mga2</i>	CTTATCCAACATTCTCAGAAATCC	CGTCGAGATGTAACCCAAATTTG	72
<i>msmR</i>	TTTGTCACTACATTATTGCCATTG	AGCTCTTCTTTGCTACGATGAG	117
<i>nra</i>	AACGTCATCACAACTCTGA	TCTCAAGCCAAATTCCCCTAG	112
<i>prtF2</i>	AATATGCCTCTAAGTATACAAG	TTAGTTGGGTAACCTTCTGTC	96
<i>recA</i>	ATTGATTGATTCTGGTGCGG	ATTTACGCATGGCCTGACTC	139
<i>sipA2</i>	TGAGTCCTGCTTTAAGTGCAAGTG	AGCTTGAGCGACAATTCGACCAAC	127
<i>ska</i>	CGCAATGCCACATAAACTTG	AAGTAGACCTTGCCGTTTCG	130
<i>speB</i>	TGTCGGTAAAGTAGGCGGAC	GAGCTGAAGGGTTTAGTGCG	130
<i>srtC2</i>	GTTAGCTTTGAATGAGGATGTC	ATGGCAACACTGCCATCAACTG	125
<i>up-3</i>	GGTTGCAGGAACTAATATTAG	CCCTCCTTCTAAACTAAAGTGGC	132
<i>up-4</i>	GAAACAGGCGAATGATTGACC	CAGTAATCCGATCGTCGTTTGTGCGCG	139