Table S9. Lambda chain family usage of isolated influenza-specific rmAbs.

| | Influenza-Specific rmAbs | | | | | | | | | | | |
|---------|--------------------------|------------|------------|----|---|----------|----------|---|---|----------|--------------|-------|
| Subject | Lambda Chain Family | | | | | | | | | | | |
| | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | Total |
| | | | | | | N (%) | | | | | | N |
| TIV01 | 33 (47.1%) | 10 (14.3%) | 23 (32.9%) | _* | _ | 1 (1.4%) | 2 (2.9%) | - | - | 1 (1.4%) | - | 70 |
| TIV04 | 5 (62.5%) | 1 (12.5%) | 2 (25%) | - | - | - | - | - | - | - | - | 8 |
| TIV14 | 1 (100%) | - | - | - | - | - | - | - | - | - | - | 1 |
| TIV21 | 3 (20%) | 2 (13.3%) | 9 (60%) | - | - | 1 (6.7%) | - | - | - | - | - | 15 |
| TIV24 | 9 (64.3%) | - | 4 (28.6%) | - | = | - | - | - | - | 1 (7.1%) | - | 14 |
| total | 51 (47.2%) | 13 (12.0%) | 38 (35.2%) | - | - | 2 (1.9%) | 2 (1.9%) | - | - | 2 (1.9%) | - | 108 |
| | | | | | | | | | | | | |
| EI02 | 1 (100%) | - | - | - | - | - | - | - | - | - | - | 1 |
| EI03 | 2 (100%) | - | - | - | - | - | - | - | - | - | - | 2 |
| EI05 | 1 (100%) | - | - | - | - | - | - | - | - | - | - | 1 |
| EI07 | - | - | - | - | - | - | = | - | - | - | - | 0 |
| EI12 | - | - | - | - | - | - | - | - | - | - | - | 0 |
| EI13 | 4 (80%) | 1 (20%) | - | - | - | - | = | = | = | - | - | 5 |
| total | 8 (88.9%) | 1 (11.1%) | - | - | - | - | - | - | - | - | - | 9 |

^{* - =} No antibodies of this lambda chain family isolated.