

Psychotropic medication group	AN (n=54) <i>n (%)</i>	BN (n=585) <i>n (%)</i>	EDNOS-P (n=61) <i>n (%)</i>	BED (n=3104) <i>n (%)</i>	No ED (n=58215) <i>n (%)</i>
Antidepressants					
Before pregnancy	9 (16.7)[*]	46 (7.9)[*]	6 (9.8)[‡]	156 (5.0)[*]	1474 (2.5)
First trimester	5 (9.3)[*]	24 (4.1)[*]	5 (8.2)[*]	66 (2.1)[*]	469 (0.8)
Second trimester	3 (5.6)[‡]	15 (2.6)[*]	3 (4.9)[‡]	33 (1.1)[*]	254 (0.4)
Third trimester	1 (1.9)	9 (1.5)[*]	2 (3.3)	28 (0.9)[*]	185 (0.3)
Any time during pregnancy	7 (13.0)[*]	33 (5.6)[*]	5 (8.2)[*]	86 (2.8)[*]	586 (1.0)
0-3 months postpartum	-	9 (1.5)[*]	1 (1.6)	25 (0.8)[*]	236 (0.4)
4-6 months postpartum	2 (3.7)	13 (2.2)[*]	2 (3.3)	35 (1.1)[*]	359 (0.6)
Before, during and after pregnancy	1 (1.9)	7 (1.2)[*]	1 (1.6)	17 (0.5)[‡]	159 (0.3)
Anxiolytics and sedatives					
Before pregnancy	5 (9.3)[*]	21 (3.6)[*]	1 (1.6)	34 (1.1)	572 (1.0)
First trimester	1 (1.9)	11 (1.9)[*]	-	16 (0.5)	257 (0.4)
Second trimester	-	8 (1.4)[‡]	1 (1.6)	17 (0.5)	207 (0.4)
Third trimester	-	6 (1.0)[‡]	1 (1.6)	19 (0.6)[*]	160 (0.3)
Any time during pregnancy	1 (1.9)	23 (3.9)[*]	2 (3.3)	46 (1.5)[*]	494 (0.8)
0-3 months postpartum	1 (1.9)	2 (0.3)	1 (1.6)	20 (0.6)	218 (0.4)
4-6 months postpartum	3 (5.6)[‡]	7 (1.2)	3 (4.9)[‡]	22 (0.7)	257 (0.4)
Before, during and after pregnancy	1 (1.9)	-	-	3 (0.1)	44 (0.1)
Antipsychotics					
Before pregnancy	1 (1.9)	5 (0.9)[‡]	1 (1.6)	9 (0.3)	137 (0.2)
First trimester	1 (1.9)	9 (1.5)[‡]	-	19 (0.6)	344 (0.6)
Second trimester	-	7 (1.2)	-	13 (0.4)	294 (0.5)

Third trimester	1 (1.9)	1 (0.2)	-	4 (0.1)	79 (0.1)
Any time during pregnancy	2 (3.7)	12 (2.1)[‡]	-	24 (0.8)	481 (0.8)
0-3 months postpartum	1 (1.9)	1 (0.2)	-	4 (0.1)	51 (0.1)
4-6 months postpartum	1 (1.9)	2 (0.3)	-	3 (0.1)	60 (0.1)
Before, during and after pregnancy	-	1 (0.2)	-	-	18 (0.03)
