S4 Table. The effect of depression on the occurrence of chronic pain and/or angina in the "unrelated" subgroup (n=9,163) and stratified according to gender.

Exposure	Outcome	Group	Unadjusted		Adjusted	
_		_	N	OR [95% CI]	N	OR [95% CI]
Depression in the presence of chronic pain						
Chronic Pain	Depression	Overall [†]	4,780	2.70 a	4,179	2.46 a, b
emonic rum	Depression	Overan	1,700	[2.26 to 3.23]	1,17	[2.02 to 3.01]
		Females	2,743	2.64 ^a	2,385	2.74 a,c
		only [‡]		[2.13 to 3.26]		[2.16 to 3.46]
		Males only [‡]	2,037	2.25 a	1,794	2.05 a, d
				[1.59 to 3.17]		[1.41 to 2.98]
Depression in the presence of angina						
Angina	Depression	Overall [†]	6,264	2.08 a	5,453	2.08 a
-				[1.73 to 2.51]		[1.67 to 2.60]
		Females	3,662	2.08 a	3,165	2.07 a, e
		only [‡]		[1.66 to 2.62]		[1.59 to 2.69]
		Males only [‡]	2,602	2.28 a	2,288	2.25 a, f
				[1.63 to 3.21]		[1.52 to 3.33]
Depression in the presence of chronic pain and angina						
Chronic pain	Depression	Overall [†]	3,149	4.33 a	2,769	4.23 a, g
and angina				[3.23 to 5.81]		[3.00 to 5.96]
		Females	1,783	4.10 a	1,566	4.77 a, h
		only [‡]		[2.90 to 5.80]		[3.23 to 7.05]
		Males only‡	1,366	4.12 a	1,203	3.96 ^{a, i}
				[2.34 to 7.27]		[2.05 to 7.66]

†valid data adjusted for age, gender, education, SIMD and smoking status; ‡valid data adjusted for age, education, SIMD and smoking status; a= p<0.001; b= Variables not in final model: Education (P=0.14); c= Variables not in final model: Education (P=0.38); SIMD (P=0.21); d= Variables not in final model: Education (P=0.20); SIMD (P=0.48); e= Variables not in final model: SIMD (P=0.30); f= Variables not in final model: Education (P=0.40); g= Variables not in final model: SIMD (P=0.12); Education (P=0.28); h= Variables not in final model: Smoking (P=0.10); SIMD (P=0.34); Education (P=0.22); i= Variables not in final model: SIMD (P=0.17); Education (P=0.94)